## Homework 1, due Friday, August 24

Please turn in well-written solutions for the following problems:

- (1) (1.1.13 in Tao) Let X be any set,  $x \in X$ , and  $(x_n)_{n=1}^{\infty}$  a sequence in X. Prove that  $(x_n)$  converges to x in the discrete metric  $d_{\text{disc}}$  if and only if there exists N such that  $x_n = x$  for every  $n \ge N$ .
- (2) (1.1.16 in Tao) Let  $(x_n)_{n=1}^{\infty}$  and  $(y_n)_{n=1}^{\infty}$  be two sequences in some metric space (X,d), such that  $x_n \to x$  and  $y_n \to y$  for some points  $x,y \in X$ . Prove that  $\lim_{n\to\infty} d(x_n,y_n) = d(x,y)$ . (Hint: Use the triangle inequality multiple times.)
- (3) Let X be the set of all continuous real-valued functions with domain [0,1]. Define a function  $d: X \times X \to [0,\infty)$  by  $d(f,g) = \int_0^1 (f(x) g(x))^2 dx$ , for any f and g in X. Prove that (X,d) is NOT a metric space, because the triangle inequality is not satisfied. (Hint: Consider constant functions.)
- (4) Consider  $\mathbb{R}^2$  with the metrics  $d_{l^2}$ ,  $d_{l^1}$ ,  $d_{l^{\infty}}$ , and  $d_{\text{disc}}$ . In each of these metrics, sketch B((0,0),1), the ball of radius 1 centered at the origin. That is, I want you to:
  - (i) Sketch  $B_{(\mathbb{R}^2,d_{12})}((0,0),1)$ .
  - (ii) Sketch  $B_{(\mathbb{R}^2,d_{l^1})}((0,0),1)$ .
  - (iii) Sketch  $B_{(\mathbb{R}^2, d_{1\infty})}((0,0), 1)$ .
  - (iv) Sketch  $B_{(\mathbb{R}^2, d_{\text{disc}})}((0, 0), 1)$ .
- (5) Let (X, d) be a metric space, and let  $E, F \subset X$ . Recall the notation that  $A \setminus B$  is the set of all elements in A that are not elements in B. Using the definitions and theorems given in class, prove the following:
  - (a)  $int(E) = E \setminus \partial E$
  - (b)  $int(E) \cap int(F) = int(E \cap F)$
  - (c)  $\operatorname{int}(E) \cup \operatorname{int}(F) \subset \operatorname{int}(E \cup F)$  (also, show by example that equality is not always true!)
- (6) Let (X,d) be a metric space, and let  $E \subset X$ . Prove that  $x \in \partial E$  if and only if for every r > 0,  $B(x,r) \cap E \neq \emptyset$  and  $B(x,r) \cap E^c \neq \emptyset$ .

In addition, I suggest that you study these problems from Tao:

- Section 1.1, problems 1.1.4, 1.1.5, 1.1.6, 1.1.12
- Section 1.2, problems 1.2.1, 1.2.4