

On nowhere weakly symmetric functions and functions with two element range

Krzysztof Ciesielski

Department of Mathematics, West Virginia University,

Morgantown, WV 26506-6310, USA

e-mail: K.Cies@math.wvu.edu

web page: <http://www.math.wvu.edu/~kcies>

Kandasamy Muthuvel

Department of Mathematics, University of Wisconsin-Oshkosh

Oshkosh, Wisconsin 54901-8601

e-mail: muthuvel@vaxa.cis.uwosh.edu

Andrzej Nowik

Current address: Institute of Mathematics Polish Academy of Sciences

Abrahama 18, Sopot, Poland

Permanent address: Department of Mathematics, Gdańsk University

ul. Wita Stwosza 57, 80-952 Gdańsk, Poland

e-mail: matan@paula.univ.gda.pl

web page: <http://math.univ.gda.pl/~tm/a.nowik>

Abstract

A function $f: \mathbb{R} \rightarrow \{0, 1\}$ is weakly symmetric (weakly symmetrically continuous) at $x \in \mathbb{R}$ provided there is a sequence $h_n \rightarrow 0$ such

AMS classification numbers: Primary 26A15; Secondary 03E50, 03E35.

Key words and phrases: continuum hypothesis, second category sets, two-element range functions, symmetric functions, symmetrically continuous functions, anti-Schwartz functions.

that $f(x + h_n) = f(x - h_n) = f(x)$ ($f(x + h_n) = f(x - h_n)$, respectively) for every n . We will characterize the sets $S(f)$ of all points at which f fails to be weakly symmetrically continuous and show that f must be weakly symmetric at some $x \in \mathbb{R} \setminus S(f)$. In particular, there is no $f: \mathbb{R} \rightarrow \{0, 1\}$ which is nowhere weakly symmetric.

It is also shown that if at each point x we ignore some countable set from which we can choose the sequence h_n , then there exists a function $f: \mathbb{R} \rightarrow \{0, 1\}$ which is nowhere weakly symmetric in this weaker sense if and only if the continuum hypothesis holds.

1 Introduction

The terminology of this note is standard and follows [?].

Recall that a function $f: \mathbb{R} \rightarrow \mathbb{R}$ is *symmetrically continuous* provided

$$\lim_{h \rightarrow 0} [f(x + h) - f(x - h)] = 0 \tag{1}$$

for every $x \in \mathbb{R}$. A “total negation” of the condition (??) leads to a notion of *nowhere weakly symmetrically continuous* function,¹ that is,

$$\lim_{n \rightarrow \infty} [f(x + h_n) - f(x - h_n)] \neq 0 \tag{2}$$

for every sequence $h_n \searrow 0$. It is easy to see that if the range of f is discrete then (??) is equivalent to: “ $f(x + h_n) \neq f(x - h_n)$ for all but finitely many n .” In particular, if $f: \mathbb{R} \rightarrow \mathbb{N}$ then f is not weakly symmetrically continuous at $x \in \mathbb{R}$ provided $f(x + h) \neq f(x - h)$ for all $h > 0$ small enough, that is, when there exists an $\varepsilon_x > 0$ such that the interval $(0, \varepsilon_x)$ is disjoint with the set

$$S_x = \{h > 0: f(x - h) = f(x + h)\}.$$

In [?] Ciesielski and Larson constructed a nowhere weakly symmetrically continuous function $f: \mathbb{R} \rightarrow \mathbb{N}$ for which each set S_x is finite. (See also [?, cor. 7.4.2].) They show also there that there is no nowhere weakly symmetrically continuous function with two-element range. The latter fact was proved earlier by Kostyrko [?].

¹This terminology comes from a monograph Thomson [?]. Ciesielski and Larson [?] and most other papers on this subject call such a function *uniformly antisymmetric*.

Nowhere weakly symmetrically continuous functions have been studied by many authors. Komjáth and Shelah [?] investigated under what set theoretical assumptions the sizes of all sets S_x can be bounded by a fixed number. (See also [?].) In [?] Ciesielski proved that there is no nowhere weakly symmetrically continuous function with three-element range and in [?] he showed that the technique from the paper [?] cannot be used to prove that there is no nowhere weakly symmetrically continuous function with four-element range. It is still an open question (see [?, Problem 1(a)], [?, Problem 2(a)], or [?]) whether there exists a nowhere weakly symmetrically continuous function with finite range, though Ciesielski and Shelah [?] constructed a nowhere weakly symmetrically continuous function with bounded countable range.

For $f: \mathbb{R} \rightarrow \mathbb{R}$ let $S(f)$ denote the set of all points at which f is not weakly symmetrically continuous, that is,

$$S(f) = \{x \in \mathbb{R}: f \text{ is not weakly symmetrically continuous at } x\}.$$

We say that a set $A \subset \mathbb{R}$ is locally symmetric provided for every $x \in A$ there exists an $\varepsilon_x > 0$ such that

$$x - h \in A \iff x + h \in A \quad \text{for every } h \in (0, \varepsilon_x).$$

Our first theorem characterizes the family of all sets $S(f)$ with $f: \mathbb{R} \rightarrow \{0, 1\}$.

Theorem 1 (a) *If $f: \mathbb{R} \rightarrow \{0, 1\}$ then the set $S(f)$ is countable and locally symmetric.*

(b) *For every countable and locally symmetric subset A of \mathbb{R} there exists an $f: \mathbb{R} \rightarrow \{0, 1\}$ such that $S(f) = A$.*

A notion dual to weakly symmetrically continuous function is that of a symmetric function: a function $f: \mathbb{R} \rightarrow \mathbb{R}$ is a *symmetric function* provided

$$\lim_{h \rightarrow 0} [f(x+h) + f(x-h) - 2f(x)] = 0 \tag{3}$$

for every $x \in \mathbb{R}$. The “total negation” of (??) parallel to that done for the case of symmetrically continuous function leads to a notion of *nowhere weakly symmetric* function and was introduced by Ciesielski in [?].² We will

²Ciesielski [?] calls such function *anti-Schwartz*, while current terminology follows Thomson [?].

be interested in this notion only for functions $f: \mathbb{R} \rightarrow S$ when $S \subset \mathbb{N}$ contains no arithmetic progression of length three. In this particular case it is more convenient to adopt a simpler equivalent definition: a function $f: \mathbb{R} \rightarrow S$ is nowhere weakly symmetric provided for every $x \in \mathbb{R}$ there exists an $\varepsilon_x > 0$ such that the interval $(0, \varepsilon_x)$ is disjoint with the set

$$T_x = \{h > 0: f(x - h) = f(x) = f(x + h)\}.$$

The paper [?] contains also a construction of a nowhere weakly symmetric function $f: \mathbb{R} \rightarrow S \subset \mathbb{N}$ for which $T_x = \emptyset$ for every $x \in \mathbb{R}$. It also contains a question whether there exists a nowhere weakly symmetric function with two-element range. One of the goals of this paper is to answer this question in the negative:

Theorem 2 *There is no nowhere weakly symmetric function $f: \mathbb{R} \rightarrow \{0, 1\}$.*

In the second part of the paper we will consider the variations of weak symmetric continuity and weak symmetry associated with any ideal \mathcal{I} on \mathbb{R} in the following way: for every $x \in \mathbb{R}$ we choose an exceptional set $E_x \in \mathcal{I}$, and for this x we consider only the sequences h_n disjoint with E_x . The investigation of these notions for the ideal of countable subsets of \mathbb{R} leads to the following theorem.

Theorem 3 *The following conditions are equivalent.*

- (i) *The Continuum Hypothesis.*
- (ii) *There exists a function $f: \mathbb{R} \rightarrow \{0, 1\}$ such that $|S_x| \leq \omega$ for every $x \in \mathbb{R}$.*
- (iii) *There exists a function $f: \mathbb{R} \rightarrow \{0, 1\}$ such that for every $x \in \mathbb{R}$ there exists an $\varepsilon_x > 0$ such that $|S_x \cap (0, \varepsilon_x)| \leq \omega$.*
- (iv) *There exists a function $f: \mathbb{R} \rightarrow \{0, 1\}$ such that $|T_x| \leq \omega$ for every $x \in \mathbb{R}$.*
- (v) *There exists a function $f: \mathbb{R} \rightarrow \{0, 1\}$ such that for every $x \in \mathbb{R}$ there exists an $\varepsilon_x > 0$ such that $|T_x \cap (0, \varepsilon_x)| \leq \omega$.*

It should be pointed here that the equivalences (i) \Leftrightarrow (ii) \Leftrightarrow (iii) are known and were proved in [?]. (See also [?, thm. 7.4.4].)

We also show (Theorems ?? and ??) that for the ideal of meager subsets of \mathbb{R} a similar theorem cannot be proved.

2 Every function with two-element range is somewhere weakly symmetric

PROOF OF THEOREM ??(a). We will first show that $S(f)$ is countable. For $0 < n < \omega$ let

$$S_n(f) = \{x \in \mathbb{R}: (0, 1/n) \cap S_x = \emptyset\}.$$

Then clearly

$$S(f) = \bigcup_{0 < n < \omega} S_n(f).$$

We will show that each $S_n(f)$ is discrete, so countable.

Fix an $n < \omega$, $n > 0$. We will show that if $a, b \in S_n(f)$ are such that $0 < b - a < 1/n$ then the points a and b are isolated in $S_n(f)$. This clearly implies that $S_n(f)$ is discrete.

So take $a, b \in S_n(f)$ such that $0 < b - a < 1/n$ and consider the set

$$Z = \left(b - \frac{1}{n}, a + \frac{1}{n}\right) \setminus \{a, b\}.$$

We will show that $Z \cap S_n(f) = \emptyset$, so that a and b are isolated in $S_n(f)$.

Thus, take any $z \in Z$. Since the diameter of $\{a, b, z\}$ is less than $1/n$ we have

$$f(2a - z) = 1 - f(z) = f(2b - z).$$

But the mid point $(a + b) - z$ of $2a - z$ and $2b - z$ is of distance less than $1/n$ from $\{2a - z, 2b - z\}$. So $(a + b) - z \notin S_n(f)$.

This implies that the set $Z = (a + b) - Z$ is disjoint with $S_n(f)$, as required.

It is also easy to see that if $x \in S(f)$ and $\varepsilon_x > 0$ is such that $(0, \varepsilon_x) \cap S_x = \emptyset$ then

$$x - h \in S(f) \iff x + h \in S(f) \quad \text{for every } h \in (0, \varepsilon_x).$$

Thus, $S(f)$ is locally symmetric. □

PROOF OF THEOREM ??(b). The proof presented here is an elaboration of that for [?, thm. 1], in which the authors prove that for any countable subset A of \mathbb{R} there exists an $f: \mathbb{R} \rightarrow \{0, 1\}$ with $A \subset S(f)$.

Let $\{a_n: 0 < n < \omega\}$ be an enumeration of A and $\{d_n: 0 < n < \omega\}$ be an enumeration of $G(A) \setminus A$, where $G(A)$ is an additive group generated by A . Following [?] for every $n < \omega$ we define inductively a finite collection

$\mathcal{I}_n = \{I_\gamma: \gamma \in \Gamma_n\}$ of open intervals such that $\emptyset = \Gamma_0 \subseteq \Gamma_1 \subseteq \dots$, each I_γ is of the form $(b_\gamma - h_\gamma, b_\gamma + h_\gamma)$, and the following properties hold for any $\gamma, \gamma' \in \Gamma_n$.

- (1) If $\gamma \neq \gamma'$, then either $I_\gamma \cap I_{\gamma'} = \emptyset$ or one of the intervals I_γ and $I_{\gamma'}$ is a subset of another.
- (2) If $I_{\gamma'} \subseteq I_\gamma$, then either $I_{\gamma'} \subseteq (b_\gamma - h_\gamma, b_\gamma)$ or $I_{\gamma'} \subseteq (b_\gamma, b_\gamma + h_\gamma)$.
- (3) $\{a_1, \dots, a_n\} \subseteq B_n \subseteq A$, where $B_n = \{b_\gamma: \gamma \in \Gamma_n\}$.
- (4) $b_\gamma \pm h_\gamma \notin A$.
- (5) If $I_{\gamma'} \subseteq I_\gamma$, then $\varphi_\gamma[I_{\gamma'}] \in \mathcal{I}_n$, where $\varphi_\gamma: I_\gamma \rightarrow I_\gamma$ is a symmetry function with respect to b_γ , that is, $\varphi_\gamma(x) = 2b_\gamma - x$.
- (6) If $x \in I_\gamma$ then $x \in A$ if and only if $\varphi_\gamma(x) \in A$.
- (7) If $I_\gamma \notin \mathcal{I}_{n-1}$ then $d_n \notin I_\gamma$.

Conditions (1)–(5) are present in the proof of [?, thm. 1], while (6) and (7) are specific for our construction.

The construction is pretty much repetition of that in [?]. We start with $\Gamma_0 = \mathcal{I}_0 = \emptyset$. If \mathcal{I}_{n-1} is already constructed, we construct \mathcal{I}_n as follows. If $a_n \in B_{n-1}$ we put $\mathcal{I}_n = \mathcal{I}_{n-1}$. So assume that $a_n \notin B_{n-1}$ and notice that the set B of all (well defined) numbers $\varphi_{\gamma_1} \circ \dots \circ \varphi_{\gamma_r}(a_n)$ has at most $2^{|\mathcal{I}_{n-1}|}$ elements and that $a_n \in B \subset A \setminus B_{n-1}$. Find an $h > 0$ small enough that for every $b \in B$ the interval $(b - h, b + h)$ is disjoint with the set $\{d_1, \dots, d_n\} \cup B_n \cup \{b_\gamma \pm h_\gamma: \gamma \in \Gamma_{n-1}\}$ and that $A \cap (b - h, b + h)$ is symmetric with respect to b . Moreover, decreasing h if necessary, we also require that $\{b \pm h: b \in B\}$ is disjoint with A . We define \mathcal{I}_n as $\mathcal{I}_{n-1} \cup \{(b - h, b + h): b \in B\}$. It is easy to see that \mathcal{I}_n is as desired.

In what follows we will write $y = \varphi_{\gamma_1} \circ \dots \circ \varphi_{\gamma_r}(x)$ only when all γ_i 's are from $\Gamma = \bigcup_{n < \omega} \Gamma_n$ and the value of the composition is well defined. Note also that if $y = \varphi_{\gamma_1} \circ \varphi_{\gamma_2}(x)$ and $I_{\gamma_1} \subsetneq I_{\gamma_2}$ then we have also $y = \varphi_{\gamma_2} \circ \varphi_{\gamma'_1}(x)$, where $b_{\gamma'_1} = \varphi_{\gamma_2}(b_{\gamma_1})$. From this and the fact that $\varphi_\gamma \circ \varphi_\gamma(z) = z$ for every z we can conclude that

- (*) If $y = \varphi_{\gamma'_1} \circ \dots \circ \varphi_{\gamma'_s}(x)$ there exists an $r \leq s$ of the same parity that s and $\gamma_1, \dots, \gamma_r \in \Gamma$ such that $y = \varphi_{\gamma_1} \circ \dots \circ \varphi_{\gamma_r}(x)$ and $I_{\gamma_1} \supsetneq \dots \supsetneq I_{\gamma_r}$.

Notice also the following two facts.

(A) If $y = \varphi_{\gamma_1} \circ \cdots \circ \varphi_{\gamma_r}(x)$ and r is odd then $y \neq x$.

Indeed, by (*) we can assume that $I_{\gamma_1} \supsetneq \cdots \supsetneq I_{\gamma_r}$. But this means that x and y are on the different sides of b_{γ_1} , so they are not equal.

(B) For every $x_0 \in \mathbb{R} \setminus A$ there exists a $\delta = \delta_x > 0$ such that for every $x \in (x_0 - \delta, x_0 + \delta)$ if $y = \varphi_{\gamma_1} \circ \cdots \circ \varphi_{\gamma_r}(x)$ and r is odd then $y \neq 2x_0 - x$.

To see it, first notice that

$$y = \varphi_{\gamma_1} \circ \cdots \circ \varphi_{\gamma_r}(x) = -x - 2 \sum_{i=1}^r (-1)^i b_{q_i}. \quad (4)$$

Now, if $2x_0 - x = y = \varphi_{\gamma_1} \circ \cdots \circ \varphi_{\gamma_r}(x)$ then $x_0 = -\sum_{i=1}^r (-1)^i b_{q_i} \in G(A)$. In particular, if $x_0 \notin G(A)$ then (B) holds for any $\delta > 0$. Thus, we can assume that $x_0 \in G(A) \setminus A$, that is, $x_0 = d_n$ for some $n < \omega$. Note also that if x_0 does not belong to any I_γ then y stays on the same side of x_0 that x does and, once again, any $\delta > 0$ works. So we can assume that x_0 belongs to some I_γ which, by (7), must belong to \mathcal{I}_{n-1} .

Let I_γ be the unique shortest interval such that $x_0 \in I_\gamma$ and let $\delta > 0$ be such that $(x_0 - \delta, x_0 + \delta) \subset (b_\gamma - h_\gamma, b_\gamma + h_\gamma) \setminus \{b_\gamma\}$. We will show that this δ satisfies (B).

Assume, by way of contradiction, that there is an $x \in (x_0 - \delta, x_0 + \delta)$ such that $2x_0 - x = y = \varphi_{\gamma_1} \circ \cdots \circ \varphi_{\gamma_r}(x)$ and r is odd. By (*) we can also assume that $I_{\gamma_1} \supsetneq \cdots \supsetneq I_{\gamma_r}$.

Since b_{q_1} stays on one side of each I_{q_i} , $1 < i \leq r$, the points x and $\varphi_{\gamma_2} \circ \cdots \circ \varphi_{\gamma_r}(x)$ are on the same side of b_{q_1} . So x and $2x_0 - x = \varphi_{\gamma_1} \circ \cdots \circ \varphi_{\gamma_r}(x)$ are on the opposite sides of b_{q_1} . But $x, 2x_0 - x \in (x_0 - \delta, x_0 + \delta) \subset I_\gamma$, implying that $b_{q_1} \in (x_0 - \delta, x_0 + \delta) \subset I_\gamma$. So, $I_{\gamma_1} \subsetneq I_\gamma$. However, x and $2x_0 - x = y$ belong also to I_{γ_1} , and so does x_0 , which is between x and $2x_0 - x$. This gives $x_0 \in I_{\gamma_1} \subsetneq I_\gamma$, contradicting the choice of I_γ . This completes the proof of (B).

With (A) and (B) in hand, we can construct the desired function f . For this, consider \mathbb{R} as vertices of an infinite graph and declare numbers x and y as connected by an edge provided $y = \varphi_\gamma(x)$ for some $\gamma \in \Gamma$. Also, for each $x \in \mathbb{R}$ let C_x stands for all $y \in \mathbb{R}$ such that there exists a path from x to y , that is, such that $y = \varphi_{\gamma_1} \circ \cdots \circ \varphi_{\gamma_r}(x)$ for some $\gamma_1, \dots, \gamma_r \in \Gamma$.

Note that condition (A) says that our graph has no odd cycles, so we can color it by two colors, 0 and 1, such that no connected vertices have the same color. In fact, if we color each component C_x that way, then the resulted coloring, a function $f: \mathbb{R} \rightarrow \{0, 1\}$, will have this property. It is easy to see that for any such f we will have $A \subset S(f)$. To see that we can find such an f with $A = S(f)$ first note that, by (??), if $y \in C_x$ then y belongs to the group generated by $A \cup \{x\}$. Thus each component C_x is at most countable. We color components C_x by transfinite induction. We enumerate $(\mathbb{R} \setminus A) \times \omega$ as $\langle (x_\xi, n_\xi) : \xi < \mathfrak{c} \rangle$. Next, at each stage $\xi < \mathfrak{c}$ we choose and color two new components C_ξ and C'_ξ in the following way. First we choose $x \in (x_\xi, x_\xi + \delta_{x_\xi}/2^{n_\xi})$ such that neither x nor $2x_\xi - x$ is in previously chosen components. Then we choose components C_ξ and C'_ξ containing x and $2x_\xi - x$, respectively, and color each in such a way, that both x and $2x_\xi - x$ get the same color. Such a coloring can be obviously chosen if components C_ξ and C'_ξ are not equal. However, if $C_\xi = C'_\xi$ then, by (B), x and $2x_\xi - x$ must be colored the same color automatically.

It is easy to see that a coloring function f defined that way is as desired.

□

The characterization given in Theorem ?? does not hold for functions with a 3-element range. This follows from the fact that, by [?, thm. 2], for every Hamel basis $H \subset \mathbb{R}$ there exists an $f: \mathbb{R} \rightarrow \{0, 1, 2\}$ such that $H \subset S(f)$. Since a Hamel basis can be a Bernstein set (see e.g. [?, cor. 7.3.7]) the set $S(f)$ for such an f need to be neither meager nor of measure zero.

With Theorem ?? in hand we are ready to prove our second theorem.

PROOF OF THEOREM ??. By way of contradiction assume that f is nowhere weakly symmetric.

From Theorem ?? we know that $|S(f)| \leq \omega$. Translating f , if necessary, we can assume that $0 \notin S(f)$. Let V be an uncountable linear subspace of \mathbb{R} over \mathbb{Q} disjoint with $S(f)$. We will obtain a contradiction with [?, thm. 2.1] by showing that the restriction function

$$f \upharpoonright V: V \rightarrow \{0, 1\}$$

is nowhere weakly symmetrically continuous.

So, fix an $x \in V$. Since f is not weakly symmetric at x there exists an $\varepsilon > 0$ such that $(0, \varepsilon) \cap T_x = \emptyset$. We will show that this implies that we have also $(0, \varepsilon) \cap S_x \cap V = \emptyset$. We will assume that

$$f(x) = 0,$$

the argument for $f(x) = 1$ being essentially the same.

By way of contradiction assume that there is an $h \in (0, \varepsilon) \cap S_x \cap V$. Then $f(x - h) = f(x + h)$ is not equal to $f(x) = 0$ since $h \notin T_x$. In particular,

$$f(x - h) = f(x + h) = 1.$$

Since f is not weakly symmetric at $x+h$ and at $x-h$ we can find a $\delta \in (0, \varepsilon-h)$ such that $(T_{x-h} \cup T_{x+h}) \cap (0, \delta) = \emptyset$.

Since $x - h \in V \subset \mathbb{R} \setminus S(f)$ there exists a $t \in (0, \delta) \cap (0, h) \cap S_{x-h}$. Then $f((x - h) - t) = f((x - h) + t) \neq f(x - h) = 1$, since $t \in S_{x-h} \setminus T_{x-h}$. Thus,

$$f((x - h) - t) = f((x - h) + t) = 0.$$

Since $h + t \in (0, \varepsilon) \subset \mathbb{R} \setminus T_x$ and $f((x - h) - t) = f(x) = 0$ we conclude that

$$f((x + h) + t) = 1.$$

Similarly $h - t \in (0, \varepsilon) \subset \mathbb{R} \setminus T_x$ and $f((x - h) + t) = f(x) = 0$ imply that

$$f((x + h) - t) = 1.$$

Thus $f((x + h) + t) = f((x + h) - t) = f(x + h) = 1$ contradicting the fact that $t \notin T_{x+h}$. \square

Now for $f: \mathbb{R} \rightarrow \mathbb{R}$ let $T(f)$ be the set of all $x \in \mathbb{R}$ such that f is not weakly symmetric at x . In this notation Theorem ?? says that $T(f) \neq \mathbb{R}$ for every function $f: \mathbb{R} \rightarrow \{0, 1\}$. One might suspect that a characterization similar to that from Theorem ?? may be true also for $T(f)$. This, however, is not true.

Example 4 *There exists an $f: \mathbb{R} \rightarrow \{0, 1\}$ such that $T(f)$ is not meager.*

PROOF. Let $f_0: \mathbb{R} \rightarrow \mathbb{N}$ be a function from [?, cor. 1.2]. For this function none of the level sets $f_0^{-1}(n)$ contains an arithmetic progression of length 3. Let $n \in \mathbb{N}$ be such that $f_0^{-1}(n)$ is not meager and let f be a characteristic function of the set $f_0^{-1}(n)$. It is easy to see that $f_0^{-1}(n) \subset T(f)$. \square

It is easy to see that the same argument gives also a function $f: \mathbb{R} \rightarrow \{0, 1\}$ for which the set $T(f)$ is not of measure zero.

It is also worth to mention here the following fact.

Proposition 5 (Szyszkowski [?]) *For every $B \subset \mathbb{R}$ there exists an $f: \mathbb{R} \rightarrow \mathbb{N}$ for which $S(f) = B$.*

We do not know whether its weakly symmetric analog is also true. Therefore the following question remains open.

Problem 6 *Is it true that for every $B \subset \mathbb{R}$ there exists an $f: \mathbb{R} \rightarrow \mathbb{N}$ for which $T(f) = B$?*

3 Nowhere almost weakly symmetric function with two-element range

We will start this section with the proof of Theorem ???. Since the equivalences (i) \Leftrightarrow (ii) \Leftrightarrow (iii) are known it remains to show (i) \Leftrightarrow (iv) \Leftrightarrow (v). This follows immediately from the following theorem.

Theorem 7 *Let κ be an infinite cardinal number and let W be a linear subspace of \mathbb{R} over \mathbb{Q} . The following conditions are equivalent.*

- (i) $|W| \leq \kappa^+$.
- (ii) *There exists a function $f: \mathbb{R} \rightarrow \{0, 1\}$ such that $|T_x \cap W| \leq \kappa$ for every $x \in W$.*
- (iii) *There exists a function $f: \mathbb{R} \rightarrow \{0, 1\}$ such that for every $x \in W$ there exists an $\varepsilon_x > 0$ such that $|T_x \cap W \cap (0, \varepsilon_x)| \leq \kappa$.*

We will prove Theorem ??? only for $\kappa = \omega$ which is enough to get Theorem ???. The proof of the general case is essentially the same. (Though it requires also a simple generalization of the first part of Theorem ???.) So, in the rest of the argument we will assume that $\kappa = \omega$.

Since $T_x \subset S_x$, the implication (i) \Rightarrow (ii) follows from the first part of Theorem ???. Clearly (ii) implies (iii). Thus, it is enough to prove only that (iii) implies (i).

In the proof we will need the following lemma.

Lemma 8 *Let $f: \mathbb{R} \rightarrow \{0, 1\}$ and let V be a linear subspace of \mathbb{R} over \mathbb{Q} . If $Z = \{x \in V: \exists \varepsilon > 0 (|S_x \cap V \cap (0, \varepsilon)| \leq \omega)\}$ then $|Z| \leq \omega_1$.*

PROOF. The proof is essentially identical to that for (iii) \Rightarrow (i) from Theorem ?? as presented in [?, thm. 7.4.4]. Just assume, by way of contradiction, that $|Z| > \omega_1$ and choose $B \subset Z$ of cardinality ω_2 which is linearly independent over \mathbb{Q} . For this choice of B the intact proof presented in [?, thm. 7.4.4] gives the desired contradiction. \square

PROOF OF THEOREM ?. Let f be as in (iii) and, by way of contradiction, assume that $|W| \geq \omega_2$. Let Z_0 be as in Lemma ?? with $V = W$. By shifting the domain of f , if necessary, we can assume that $0 \notin Z_0$. Let V be a linear subspace of W over \mathbb{Q} such that $|V| \geq \omega_2$ and $V \cap Z_0 = \emptyset$. Apply again Lemma ?? to get a set Z working for f and this V . Then $|V \setminus Z| \geq \omega_2$.

Fix an $x \in V \setminus Z$. By (iii) we can choose an $\varepsilon > 0$ with the property that $|T_x \cap W \cap (0, 2\varepsilon)| \leq \omega$. Since $x \notin Z$ we can also find an $h \in (0, \varepsilon) \cap S_x \cap V \setminus T_x$. In particular $f(x - h) = f(x + h)$. Next, assume that

$$f(x) = 0,$$

the argument for $f(x) = 1$ being essentially the same. Then

$$f(x - h) = f(x + h) = 1,$$

since $h \notin T_x$. Let $\delta \in (0, h)$ be such that the set $T = (T_{x-h} \cup T_{x+h}) \cap W \cap (0, \delta)$ is countable. Since $x - h \in V \subset W \setminus Z_0$, the set $S_{x-h} \cap W \cap (0, \delta)$ is uncountable. In particular, there exists a $t \in S_{x-h} \cap W \cap (0, \delta) \setminus T$ such that

$$h + t, h - t \in (0, 2\varepsilon) \setminus T_x.$$

Then $f((x - h) - t) = f((x - h) + t) \neq f(x - h) = 1$, since $t \in S_{x-h} \setminus T_{x-h}$. Thus,

$$f((x - h) - t) = f((x - h) + t) = 0.$$

Since $h + t \notin T_x$ and $f((x - h) - t) = f(x) = 0$ we conclude that

$$f((x + h) + t) = 1.$$

Similarly $h - t \notin T_x$ and $f((x - h) + t) = f(x) = 0$ imply that

$$f((x + h) - t) = 1.$$

Thus $f((x + h) + t) = f((x + h) - t) = f(x + h) = 1$ contradicting the fact that $t \notin T_{x+h}$. \square

Let us finish with a remark that Theorem ?? can be proved with W being replaced by any additive subgroup of \mathbb{R} .

The above theorems show that, under CH, for a function $f: \mathbb{R} \rightarrow \{0, 1\}$ the sets S_x and T_x can be countable. Can we make them any smaller? Can they all be, for example, nowhere dense? The following two theorems show that this is not possible. In what follows \mathcal{M}_0 and \mathcal{M} will stand for the ideals of nowhere dense and meager subsets of \mathbb{R} , respectively.

Theorem 9 *Let $f: \mathbb{R} \rightarrow \{0, 1\}$, $M \in \mathcal{M}$, and V be a linear subspace of \mathbb{R} over \mathbb{Q} . If $S_x^M = \{h \in S_x \cap V: h \notin \pm(x - M)\}$ and*

$$Z = \{x \in V: \exists \varepsilon > 0 \ S_x^M \cap (0, \varepsilon) \in \mathcal{M}_0\}$$

then $Z \in \mathcal{M}$.

PROOF. By way of contradiction assume that $Z \notin \mathcal{M}$. So, we have also $V \notin \mathcal{M}$. For every $x \in Z$ let $n_x \in \mathbb{N}$ be such that the set

$$C_x = S_x^M \cap (0, 1/n_x)$$

is nowhere dense. The sets $Z_n = \{x \in Z \setminus M: n_x = n\}$ form a countable partition of $Z \setminus M \notin \mathcal{M}$, so there exists an $n \in \mathbb{N}$ such that $Z_n \notin \mathcal{M}$. Let U be an open interval of length less than $1/n$ such that the set $L = Z_n \cap U$ is nowhere meager in U . Note that $n_x = n$ for all $x \in L$. Moreover, if

$$C_x^* = C_x \cup \pm(x - M) \in \mathcal{M}$$

then $S_x \cap V \cap (0, 1/n) \subset C_x^*$ so that

$$f(x - h) \neq f(x + h) \text{ for } x \in L \text{ and } h \in (0, 1/n) \cap V \setminus C_x^*, \quad (5)$$

and

$$|x - y| < 1/n \text{ for every } x, y \in L.$$

Let $x = t_0 \in L$ and let $\{t_m: 0 < m < \omega\}$ be a dense subset of $L \setminus (x \pm C_x^*)$. Moreover, let K be an additive subgroup generated by $\{t_m: m < \omega\}$ and choose a

$$z \in L \setminus \bigcup_{i < \omega} (K \pm C_{t_i}^*), \quad (6)$$

$z \neq x$. Moreover, choose $y = t_m \notin \{x, z\}$ such that

$$y \notin x \pm C_z. \quad (7)$$

Such a choice is possible since C_z is nowhere dense and t_m 's are dense in U . Next define

$$a = x - y + z, \quad b = x + y - z, \quad c = -x + y + z.$$

Then $(a + b)/2 = x$, $(b + c)/2 = y$, and $(c + a)/2 = z$. We will show that

$$f(a) \neq f(b), \quad f(b) \neq f(c), \quad f(c) \neq f(a), \quad (8)$$

which will give us the desired contradiction, since the points a , b , and c are distinct and f admits only two values. But (??) follows immediately from (??) and

$$|a - x| \notin C_x^*, \quad |b - y| \notin C_y^*, \quad |c - z| \notin C_z^*, \quad (9)$$

since we obviously have $|a - x|, |b - y|, |c - z| \in (0, 1/n) \cap V$. But (??) follows easily from (??) and (??):

$|a - x| \notin C_x^*$, since otherwise we would have $z - y = a - x \in \pm C_x^*$ and $z \in y \pm C_x^*$, contradicting (??).

$|b - y| \notin C_y^*$, since otherwise we would have $z - x = y - b \in \pm C_y^*$ and $z \in x \pm C_y^*$, again contradicting (??).

$|c - z| \notin C_z^*$, since otherwise point $y - x = c - z$ would belong to the set $\pm C_z^* = \pm[C_z \cup \pm(z - M)]$. However, $y - x \in \pm C_z$ implies that $y \in x \pm C_z$, contradicting (??). On the other hand if $y - x \in \pm(z - M)$, then $z \in K + M$, contradicting (??). \square

Theorem 10 *Let $f: \mathbb{R} \rightarrow \{0, 1\}$. If*

$$U = \{x \in \mathbb{R}: \exists \varepsilon > 0 T_x \cap (0, \varepsilon) \in \mathcal{M}_0\}$$

then $\mathbb{R} \setminus U \notin \mathcal{M}$.

PROOF. The proof is quite similar to that for Theorem ???. By way of contradiction assume that $U^c = \mathbb{R} \setminus U \in \mathcal{M}$. Let $Z_0 \in \mathcal{M}$ be from Theorem ??? used with $V = \mathbb{R}$ and $M = \emptyset$. Then apply again Theorem ??? with $V = \mathbb{R}$ and $M = Z_0 \cup U^c$ to get a set $Z \in \mathcal{M}$.

Next fix an $x \in \mathbb{R} \setminus (Z \cup U^c)$. We assume that

$$f(x) = 0,$$

the argument for $f(x) = 1$ being essentially the same. Since $x \in U$ we can choose an $\varepsilon > 0$ such that $T_x \cap (0, \varepsilon) \in \mathcal{M}_0$. Since $x \notin Z$ we have $S_x^M \cap (0, \varepsilon) \notin$

\mathcal{M}_0 . Thus, there exists a non-empty open interval $J_0 \subset (0, \varepsilon) \setminus T_x$ intersecting $S_x^M = S_x \setminus \pm(x - M)$. Let $h \in S_x^M \cap J_0$. Then

$$f(x - h) = f(x + h) = 1,$$

since $h \in S_x \setminus T_x$. Since $h \notin \pm(x - M)$ we have $x \pm h \in \mathbb{R} \setminus M = U \setminus Z_0$. So there exists a $\delta > 0$ such that the set $T = (T_{x-h} \cup T_{x+h}) \cap (0, \delta)$ is nowhere dense. Decreasing δ , if necessary, we can also assume that $J = (h - \delta, h + \delta) \subset J_0$. Next, since $x - h \notin Z_0$, we can find a $t \in S_{x-h} \cap (0, \delta) \setminus T$.

Then $f((x - h) - t) = f((x - h) + t) \neq f(x - h) = 1$, since $t \in S_{x-h} \setminus T_{x-h}$. Thus,

$$f((x - h) - t) = f((x - h) + t) = 0.$$

But $h \pm t \in J \subset J_0 \subset (0, \varepsilon) \setminus T_x$ and $f(x - (h \pm t)) = f(x) = 0$ imply that

$$f(x + (h \pm t)) = 1.$$

Thus $f((x + h) \pm t) = f(x + h) = 1$ contradicting the fact that $t \notin T_{x+h}$. \square

References

- [1] F. Bagemihl, *Independent sets and second category*, Order **7** (1990), 65–66.
- [2] K. Ciesielski, *On range of uniformly antisymmetric functions*, Real Anal. Exchange **19** (1993–94), 616–619.*
- [3] K. Ciesielski, *Uniformly antisymmetric functions and K_5* , Real Anal. Exchange **21** (1995–96), 147–153.*
- [4] K. Ciesielski, *Sum and difference free partitions of vector spaces*, Colloq. Math. **71**(2) (1996), 263–271.*
- [5] K. Ciesielski, *Set Theory for the Working Mathematician*, London Math. Soc. Stud. Texts **39**, Cambridge Univ. Press 1997.
- [6] K. Ciesielski, *Set theoretic real analysis*, J. Appl. Anal. **3**(2) (1997), 143–190.*

*A preprint is available in electronic form from *Set Theoretic Analysis Web Page*: <http://www.math.wvu.edu/~kcies/STA/STA.html>

- [7] K. Ciesielski, L. Larson, *Uniformly antisymmetric functions*, Real Anal. Exchange **19** (1993–94), 226–235.*
- [8] K. Ciesielski, S. Shelah, *Uniformly antisymmetric function with bounded range*, Real Anal. Exchange **24**(2) (1998–99), 615–619.*
- [9] P. Komjáth, S. Shelah, *On uniformly antisymmetric functions*, Real Anal. Exchange **19** (1993–94), 218–225.
- [10] P. Kostyrko, *There is no strongly locally antisymmetric set*, Real Anal. Exchange **17** (1991–92), 423–425.
- [11] M. Szyszkowski, *Points of weak symmetric continuity*, Real Anal. Exchange **24** (1998–99), 807–813.
- [12] B. Thomson, *Symmetric Properties of Real Functions*, Marcel Dekker, 1994.

*A preprint is available in electronic form from *Set Theoretic Analysis Web Page*:
<http://www.math.wvu.edu/~kcies/STA/STA.html>