

Problem Set 3.1, page 127

- 1 $x + y \neq y + x$ and $x + (y + z) \neq (x + y) + z$ and $(c_1 + c_2)x \neq c_1x + c_2x$.
- 2 When $c(x_1, x_2) = (cx_1, 0)$, the only broken rule is 1 times x equals x . Rules (1)-(4) for addition $x + y$ still hold since addition is not changed.
- 3 (a) cx may not be in our set: not closed under multiplication. Also no $\mathbf{0}$ and no $-x$
 (b) $c(x + y)$ is the usual $(xy)^c$, while $cx + cy$ is the usual $(x^c)(y^c)$. Those are equal. With $c = 3, x = 2, y = 1$ this is $3(2 + 1) = 8$. The zero vector is the number 1.
- 4 The zero vector in matrix space \mathbf{M} is $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$; $\frac{1}{2}A = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$ and $-A = \begin{bmatrix} -2 & 2 \\ -2 & 2 \end{bmatrix}$.
 The smallest subspace of \mathbf{M} containing the matrix A consists of all matrices cA .
- 5 (a) One possibility: The matrices cA form a subspace not containing B (b) Yes: the subspace must contain $A - B = I$ (c) Matrices whose main diagonal is all zero.
- 6 When $f(x) = x^2$ and $g(x) = 5x$, the combination $3f - 4g$ in function space is $h(x) = 3f(x) - 4g(x) = 3x^2 - 20x$.
- 7 Rule 8 is broken: If $cf(x)$ is defined to be the usual $f(cx)$ then $(c_1 + c_2)f = f((c_1 + c_2)x)$ is not generally the same as $c_1f + c_2f = f(c_1x) + f(c_2x)$.
- 8 If $(f + g)(x)$ is the usual $f(g(x))$ then $(g + f)x$ is $g(f(x))$ which is different. In Rule 2 both sides are $f(g(h(x)))$. Rule 4 is broken there might be no inverse function $f^{-1}(x)$ such that $f(f^{-1}(x)) = x$. If the inverse function exists it will be the vector $-f$.
- 9 (a) The vectors with integer components allow addition, but not multiplication by $\frac{1}{2}$
 (b) Remove the x axis from the xy plane (but leave the origin). Multiplication by any c is allowed but not all vector additions.
- 10 The only subspaces are (a) the plane with $b_1 = b_2$ (d) the linear combinations of v and w (e) the plane with $b_1 + b_2 + b_3 = 0$.
- 11 (a) All matrices $\begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix}$ (b) All matrices $\begin{bmatrix} a & a \\ 0 & 0 \end{bmatrix}$ (c) All diagonal matrices.
- 12 For the plane $x + y - 2z = 4$, the sum of $(4, 0, 0)$ and $(0, 4, 0)$ is not on the plane. (The key is that this plane does not go through $(0, 0, 0)$.)
- 13 The parallel plane \mathbf{P}_0 has the equation $x + y - 2z = 0$. Pick two points, for example $(2, 0, 1)$ and $(0, 2, 1)$, and their sum $(2, 2, 2)$ is in \mathbf{P}_0 .
- 14 (a) The subspaces of \mathbf{R}^2 are \mathbf{R}^2 itself, lines through $(0, 0)$, and $(0, 0)$ by itself (b) The subspaces of \mathbf{R}^4 are \mathbf{R}^4 itself, three-dimensional planes $n \cdot v = 0$, two-dimensional subspaces ($n_1 \cdot v = 0$ and $n_2 \cdot v = 0$), one-dimensional lines through $(0, 0, 0, 0)$, and $(0, 0, 0, 0)$ by itself.
- 15 (a) Two planes through $(0, 0, 0)$ probably intersect in a line through $(0, 0, 0)$
 (b) The plane and line probably intersect in the point $(0, 0, 0)$
 (c) If x and y are in both \mathbf{S} and \mathbf{T} , $x + y$ and cx are in both subspaces.
- 16 The smallest subspace containing a plane \mathbf{P} and a line \mathbf{L} is either \mathbf{P} (when the line \mathbf{L} is in the plane \mathbf{P}) or \mathbf{R}^3 (when \mathbf{L} is not in \mathbf{P}).
- 17 (a) The invertible matrices do not include the zero matrix, so they are not a subspace
 (b) The sum of singular matrices $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ is not singular: not a subspace.

- 18 (a) *True*: The symmetric matrices do form a subspace (b) *True*: The matrices with $A^T = -A$ do form a subspace (c) *False*: The sum of two unsymmetric matrices could be symmetric.
- 19 The column space of A is the x -axis = all vectors $(x, 0, 0)$. The column space of B is the xy plane = all vectors $(x, y, 0)$. The column space of C is the line of vectors $(x, 2x, 0)$.
- 20 (a) Elimination leads to $0 = b_2 - 2b_1$ and $0 = b_1 + b_3$ in equations 2 and 3: Solution only if $b_2 = 2b_1$ and $b_3 = -b_1$ (b) Elimination leads to $0 = b_1 + 2b_3$ in equation 3: Solution only if $b_3 = -b_1$.
- 21 A combination of the columns of C is also a combination of the columns of A . Then $C = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix}$ and $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$ have the same column space. $B = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$ has a different column space.
- 22 (a) Solution for every \mathbf{b} (b) Solvable only if $b_3 = 0$ (c) Solvable only if $b_3 = b_2$.
- 23 The extra column \mathbf{b} enlarges the column space unless \mathbf{b} is *already in* the column space.
 $[A \ \mathbf{b}] = \begin{bmatrix} 1 & 0 & \mathbf{1} \\ 0 & 0 & \mathbf{1} \end{bmatrix}$ (larger column space) $\begin{bmatrix} 1 & 0 & \mathbf{1} \\ 0 & 1 & \mathbf{1} \end{bmatrix}$ (\mathbf{b} is in column space)
 (no solution to $A\mathbf{x} = \mathbf{b}$) ($A\mathbf{x} = \mathbf{b}$ has a solution)
- 24 The column space of AB is *contained in* (possibly equal to) the column space of A . The example $B = 0$ and $A \neq 0$ is a case when $AB = 0$ has a smaller column space than A .
- 25 The solution to $A\mathbf{z} = \mathbf{b} + \mathbf{b}^*$ is $\mathbf{z} = \mathbf{x} + \mathbf{y}$. If \mathbf{b} and \mathbf{b}^* are in $C(A)$ so is $\mathbf{b} + \mathbf{b}^*$.
- 26 The column space of any invertible 5 by 5 matrix is \mathbf{R}^5 . The equation $A\mathbf{x} = \mathbf{b}$ is always solvable (by $\mathbf{x} = A^{-1}\mathbf{b}$) so every \mathbf{b} is in the column space of that invertible matrix.
- 27 (a) *False*: Vectors that are *not* in a column space don't form a subspace. (b) *True*: Only the zero matrix has $C(A) = \{\mathbf{0}\}$. (c) *True*: $C(A) = C(2A)$.
 (d) *False*: $C(A - I) \neq C(A)$ when $A = I$ or $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ (or other examples).
- 28 $A = \begin{bmatrix} 1 & 1 & \mathbf{0} \\ 1 & 0 & \mathbf{0} \\ 0 & 1 & \mathbf{0} \end{bmatrix}$ and $\begin{bmatrix} 1 & 1 & \mathbf{2} \\ 1 & 0 & \mathbf{1} \\ 0 & 1 & \mathbf{1} \end{bmatrix}$ do not have $(1, 1, 1)$ in $C(A)$. $A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ has $C(A) = \text{line}$.
- 29 When $A\mathbf{x} = \mathbf{b}$ is solvable for all \mathbf{b} , every \mathbf{b} is in the column space of A . So that space is \mathbf{R}^9 .
- 30 (a) If \mathbf{u} and \mathbf{v} are both in $S + T$, then $\mathbf{u} = s_1 + t_1$ and $\mathbf{v} = s_2 + t_2$. So $\mathbf{u} + \mathbf{v} = (s_1 + s_2) + (t_1 + t_2)$ is also in $S + T$. And so is $c\mathbf{u} = cs_1 + ct_1$: a subspace.
 (b) If S and T are different lines, then $S \cup T$ is just the two lines (*not a subspace*) but $S + T$ is the whole plane that they span.
- 31 If $S = C(A)$ and $T = C(B)$ then $S + T$ is the column space of $M = [A \ B]$.
- 32 The columns of AB are combinations of the columns of A . So all columns of $[A \ AB]$ are already in $C(A)$. But $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ has a larger column space than $A^2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$.
 For square matrices, the column space is \mathbf{R}^n when A is *invertible*.

Problem Set 3.2, page 140

- 1 (a) $U = \begin{bmatrix} 1 & 2 & 2 & 4 & 6 \\ 0 & 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$ Free variables x_2, x_4, x_5
Pivot variables x_1, x_3 (b) $U = \begin{bmatrix} 2 & 4 & 2 \\ 0 & 4 & 4 \\ 0 & 0 & 0 \end{bmatrix}$ Free x_3
Pivot x_1, x_2
- 2 (a) Free variables x_2, x_4, x_5 and solutions $(-2, 1, 0, 0, 0)$, $(0, 0, -2, 1, 0)$, $(0, 0, -3, 0, 1)$
(b) Free variable x_3 : solution $(1, -1, 1)$. Special solution for each free variable.
- 3 The complete solution to $A\mathbf{x} = \mathbf{0}$ is $(-2x_2, x_2, -2x_4 - 3x_5, x_4, x_5)$ with x_2, x_4, x_5 free. The complete solution to $B\mathbf{x} = \mathbf{0}$ is $(2x_3, -x_3, x_3)$. The nullspace contains only $\mathbf{x} = \mathbf{0}$ when there are no free variables.
- 4 $R = \begin{bmatrix} 1 & 2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$, $R = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$, R has the same nullspace as U and A .
- 5 $A = \begin{bmatrix} -1 & 3 & 5 \\ -2 & 6 & 10 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} -1 & 3 & 5 \\ 0 & 0 & 0 \end{bmatrix}$; $B = \begin{bmatrix} -1 & 3 & 5 \\ -2 & 6 & 7 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} -1 & 3 & 5 \\ 0 & 0 & -3 \end{bmatrix} = LU$.
- 6 (a) Special solutions $(3, 1, 0)$ and $(5, 0, 1)$ (b) $(3, 1, 0)$. Total of pivot and free is n .
- 7 (a) The nullspace of A in Problem 5 is the plane $-x + 3y + 5z = 0$; it contains all the vectors $(3y + 5z, y, z) = y(3, 1, 0) + z(5, 0, 1) =$ combination of special solutions.
(b) The *line* through $(3, 1, 0)$ has equations $-x + 3y + 5z = 0$ and $-2x + 6y + 7z = 0$. The special solution for the free variable x_2 is $(3, 1, 0)$.
- 8 $R = \begin{bmatrix} 1 & -3 & -5 \\ 0 & 0 & 0 \end{bmatrix}$ with $I = [1]$; $R = \begin{bmatrix} 1 & -3 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ with $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.
- 9 (a) *False*: Any singular square matrix would have free variables (b) *True*: An invertible square matrix has *no* free variables. (c) *True* (only n columns to hold pivots)
(d) *True* (only m rows to hold pivots)
- 10 (a) Impossible row 1 (b) A is invertible (c) $A =$ all ones (d) $A = 2I, R = I$.
- 11 $\begin{bmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$
- 12 $\begin{bmatrix} 1 & 1 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$. Notice the identity matrix in the pivot columns of these *reduced* row echelon forms R .
- 13 If column 4 of a 3 by 5 matrix is all zero then x_4 is a *free* variable. Its special solution is $\mathbf{x} = (0, 0, 0, 1, 0)$, because 1 will multiply that zero column to give $A\mathbf{x} = \mathbf{0}$.
- 14 If column 1 = column 5 then x_5 is a free variable. Its special solution is $(-1, 0, 0, 0, 1)$.
- 15 If a matrix has n columns and r pivots, there are $n - r$ special solutions. The nullspace contains only $\mathbf{x} = \mathbf{0}$ when $r = n$. The column space is all of \mathbf{R}^m when $r = m$. All important!

- 16** The nullspace contains only $\mathbf{x} = \mathbf{0}$ when A has 5 pivots. Also the column space is \mathbf{R}^5 , because we can solve $A\mathbf{x} = \mathbf{b}$ and every \mathbf{b} is in the column space.
- 17** $A = [1 \ -3 \ -1]$ gives the plane $x - 3y - z = 0$; y and z are free variables. The special solutions are $(3, 1, 0)$ and $(1, 0, 1)$.
- 18** Fill in **12** then **4** then **1** to get the complete solution to $x - 3y - z = 12$: $\begin{bmatrix} x \\ y \\ z \end{bmatrix} =$
- $$\begin{bmatrix} 12 \\ 0 \\ 0 \end{bmatrix} + y \begin{bmatrix} 4 \\ 1 \\ 0 \end{bmatrix} + z \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} = \mathbf{x}_{\text{particular}} + \mathbf{x}_{\text{nullspace}}.$$
- 19** If $LU\mathbf{x} = \mathbf{0}$, multiply by L^{-1} to find $U\mathbf{x} = \mathbf{0}$. Then U and LU have the same nullspace.
- 20** Column 5 is sure to have no pivot since it is a combination of earlier columns. With 4 pivots in the other columns, the special solution is $\mathbf{s} = (1, 0, 1, 0, 1)$. The nullspace contains all multiples of this vector \mathbf{s} (a line in \mathbf{R}^5).
- 21** For special solutions $(2, 2, 1, 0)$ and $(3, 1, 0, 1)$ with free variables x_3, x_4 : $R = \begin{bmatrix} 1 & 0 & -2 & -3 \\ 0 & 1 & -2 & -1 \end{bmatrix}$ and A can be any invertible 2 by 2 matrix times this R .
- 22** The nullspace of $A = \begin{bmatrix} 1 & 0 & 0 & -4 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & -2 \end{bmatrix}$ is the line through $(4, 3, 2, 1)$.
- 23** $A = \begin{bmatrix} 1 & 0 & -1/2 \\ 1 & 3 & -2 \\ 5 & 1 & -3 \end{bmatrix}$ has $(1, 1, 5)$ and $(0, 3, 1)$ in $\mathbf{C}(A)$ and $(1, 1, 2)$ in $\mathbf{N}(A)$. Which other A 's?
- 24** This construction is impossible: 2 pivot columns and 2 free variables, only 3 columns.
- 25** $A = \begin{bmatrix} 1 & -1 & 0 & 0 \\ 1 & 0 & -1 & 0 \\ 1 & 0 & 0 & -1 \end{bmatrix}$ has $(1, 1, 1)$ in $\mathbf{C}(A)$ and only the line (c, c, c, c) in $\mathbf{N}(A)$.
- 26** $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ has $\mathbf{N}(A) = \mathbf{C}(A)$ and also (a)(b)(c) are all false. Notice $\text{rref}(A^T) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$.
- 27** If nullspace = column space (with r pivots) then $n - r = r$. If $n = 3$ then $3 = 2r$ is impossible.
- 28** If A times every column of B is zero, the column space of B is contained in the nullspace of A . An example is $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$. Here $\mathbf{C}(B)$ equals $\mathbf{N}(A)$. (For $B = 0$, $\mathbf{C}(B)$ is smaller.)
- 29** For $A =$ random 3 by 3 matrix, R is almost sure to be I . For 4 by 3, R is most likely to be I with fourth row of zeros. What about a random 3 by 4 matrix?
- 31** If $\mathbf{N}(A) =$ line through $\mathbf{x} = (2, 1, 0, 1)$, A has *three pivots* (4 columns and 1 special solution). Its reduced echelon form can be $R = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$ (add any zero rows).

- 32 Any zero rows come after these rows: $R = [1 \ -2 \ -3]$, $R = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$, $R = I$.
- 33 (a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$, $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ (b) All 8 matrices are R 's!
- 34 One reason that R is the same for A and $-A$: They have the same nullspace. They also have the same column space, but that is not required for two matrices to share the same R . (R tells us the nullspace and row space.)
- 35 The nullspace of $B = [A \ A]$ contains all vectors $x = \begin{bmatrix} y \\ -y \end{bmatrix}$ for y in \mathbf{R}^4 .
- 36 If $Cx = 0$ then $Ax = 0$ and $Bx = 0$. So $N(C) = N(A) \cap N(B) = \text{intersection}$.
- 37 Currents: $y_1 - y_3 + y_4 = -y_1 + y_2 + y_5 = -y_2 + y_4 + y_6 = -y_4 - y_5 - y_6 = 0$. These equations add to $0 = 0$. Free variables y_3, y_5, y_6 : watch for flows around loops.

Problem Set 3.3, page 151

- 1 (a) and (c) are correct; (b) is completely false; (d) is false because R might have 1's in nonpivot columns.

$$2 \ A = \begin{bmatrix} 4 & 4 & 4 & 4 \\ 4 & 4 & 4 & 4 \\ 4 & 4 & 4 & 4 \\ 4 & 4 & 4 & 4 \end{bmatrix} \text{ has } R = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}. \text{ The rank is } r = 1;$$

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 5 \\ 3 & 4 & 5 & 6 \end{bmatrix} \text{ has } R = \begin{bmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}. \text{ The rank is } r = 2;$$

$$A = \begin{bmatrix} -1 & 1 & -1 & 1 \\ -1 & 1 & -1 & 1 \\ -1 & 1 & -1 & 1 \end{bmatrix} \text{ has } R = \begin{bmatrix} 1 & -1 & 1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}. \text{ The rank is } r = 1$$

$$3 \ R_A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \quad R_B = [R_A \ R_A] \quad R_C \longrightarrow \begin{bmatrix} R_A & 0 \\ 0 & R_A \end{bmatrix} \longrightarrow \begin{matrix} \text{Zero rows go} \\ \text{to the bottom} \end{matrix}$$

$$4 \ \text{If all pivot variables come last then } R = \begin{bmatrix} 0 & I \\ 0 & 0 \end{bmatrix}. \text{ The nullspace matrix is } N = \begin{bmatrix} I \\ 0 \end{bmatrix}.$$

- 5 I think $R_1 = A_1, R_2 = A_2$ is true. But $R_1 - R_2$ may have -1 's in some pivots.

- 6 A and A^T have the same rank $r =$ number of pivots. But *pivcol* (the column number)

$$\text{is 2 for this matrix } A \text{ and 1 for } A^T: A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

- 7 Special solutions in $N = [-2 \ -4 \ 1 \ 0; -3 \ -5 \ 0 \ 1]$ and $[1 \ 0 \ 0; 0 \ -2 \ 1]$.

$$8 \ \text{The new entries keep rank 1: } A = \begin{bmatrix} 1 & 2 & 4 \\ 2 & 4 & 8 \\ 4 & 8 & 16 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 6 & -3 \\ 1 & 3 & -3/2 \\ 2 & 6 & -3 \end{bmatrix},$$

$$M = \begin{bmatrix} a & b \\ c & bc/a \end{bmatrix}.$$

9 If A has rank 1, the column space is a *line* in \mathbf{R}^m . The nullspace is a *plane* in \mathbf{R}^n (given by one equation). The nullspace matrix N is n by $n - 1$ (with $n - 1$ special solutions in its columns). The column space of A^T is a *line* in \mathbf{R}^n .

$$10 \begin{bmatrix} 3 & 6 & 6 \\ 1 & 2 & 2 \\ 4 & 8 & 8 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 4 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 2 & 2 & 6 & 4 \\ -1 & -1 & -3 & -2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 3 & 2 \end{bmatrix}$$

11 A rank one matrix has one pivot. (That pivot is in row 1 after possible row exchange; it could come in any column.) The second row of U is zero.

12 Invertible r by r submatrices Use pivot rows and columns $S = \begin{bmatrix} 1 & 3 \\ 1 & 4 \end{bmatrix}$ and $S = [1]$ and $S = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.

13 P has rank r (the same as A) because elimination produces the same pivot columns.

14 The rank of R^T is also r . The example matrix A has rank 2 with invertible S :

$$P = \begin{bmatrix} 1 & 3 \\ 2 & 6 \\ 2 & 7 \end{bmatrix} \quad P^T = \begin{bmatrix} 1 & 2 & 2 \\ 3 & 6 & 7 \end{bmatrix} \quad S^T = \begin{bmatrix} 1 & 2 \\ 3 & 7 \end{bmatrix} \quad S = \begin{bmatrix} 1 & 3 \\ 2 & 7 \end{bmatrix}.$$

15 The product of rank one matrices has rank one or zero. These particular matrices have $\text{rank}(AB) = 1$; $\text{rank}(AM) = 1$ except $AM = 0$ if $c = -1/2$.

16 $(\mathbf{u}\mathbf{v}^T)(\mathbf{w}\mathbf{z}^T) = \mathbf{u}(\mathbf{v}^T\mathbf{w})\mathbf{z}^T$ has rank one unless the inner product is $\mathbf{v}^T\mathbf{w} = 0$.

17 (a) By matrix multiplication, each column of AB is A times the corresponding column of B . So if column j of B is a combination of earlier columns, then column j of AB is the same combination of earlier columns of AB . Then $\text{rank}(AB) \leq \text{rank}(B)$. No new pivot columns! (b) The rank of B is $r = 1$. Multiplying by A cannot increase this rank. The rank of AB stays the same for $A_1 = I$ and $B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$. It drops to zero for $A_2 = \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$.

18 If we know that $\text{rank}(B^T A^T) \leq \text{rank}(A^T)$, then since rank stays the same for transposes, (apologies that this fact is not yet proved), we have $\text{rank}(AB) \leq \text{rank}(A)$.

19 We are given $AB = I$ which has rank n . Then $\text{rank}(AB) \leq \text{rank}(A)$ forces $\text{rank}(A) = n$. This means that A is invertible. The right-inverse B is also a left-inverse: $BA = I$ and $B = A^{-1}$.

20 Certainly A and B have at most rank 2. Then their product AB has at most rank 2. Since BA is 3 by 3, it cannot be I even if $AB = I$.

21 (a) A and B will both have the same nullspace and row space as the R they share.
(b) A equals an *invertible* matrix times B , when they share the same R . A key fact!

$$22 A = (\text{pivot columns})(\text{nonzero rows of } R) = \begin{bmatrix} 1 & 0 \\ 1 & 4 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 4 \\ 0 & 0 & 8 \end{bmatrix}. \quad B = \begin{bmatrix} 2 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{matrix} \text{columns} \\ \text{times rows} \end{matrix} = \begin{bmatrix} 2 & 0 \\ 2 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 0 & 3 \end{bmatrix}$$

- 23 If $c = 1$, $R = \begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ has x_2, x_3, x_4 free. If $c \neq 1$, $R = \begin{bmatrix} 1 & 0 & 2 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ has x_3, x_4 free. Special solutions in $N = \begin{bmatrix} -1 & -2 & -2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ (for $c = 1$) and $N = \begin{bmatrix} -2 & -2 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$ (for $c \neq 1$). If $c = 1$, $R = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and x_1 free; if $c = 2$, $R = \begin{bmatrix} 1 & -2 \\ 0 & 0 \end{bmatrix}$ and x_2 free; $R = I$ if $c \neq 1, 2$. Special solutions in $N = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ($c = 1$) or $N = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ($c = 2$) or $N = 2$ by 0 empty matrix.
- 24 $A = [I \ I]$ has $N = \begin{bmatrix} I \\ -I \end{bmatrix}$; $B = \begin{bmatrix} I & I \\ 0 & 0 \end{bmatrix}$ has the same N ; $C = [I \ I \ I]$ has $N = \begin{bmatrix} -I & -I \\ I & 0 \\ 0 & I \end{bmatrix}$.
- 25 $A = \begin{bmatrix} 1 & 1 & 2 & 4 \\ 1 & 2 & 2 & 5 \\ 1 & 3 & 2 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} \mathbf{1} & \mathbf{0} & 2 & 3 \\ \mathbf{0} & \mathbf{1} & 0 & 1 \end{bmatrix}$ = (pivot columns) times R .
- 26 The m by n matrix Z has r ones to start its main diagonal. Otherwise Z is all zeros.
- 27 $R = \begin{bmatrix} I & F \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} r \text{ by } r & r \text{ by } n-r \\ m-r \text{ by } r & m-r \text{ by } n-r \end{bmatrix}$; $\mathbf{rref}(R^T) = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$; $\mathbf{rref}(R^T R) = \text{same } R$
- 28 The row-column reduced echelon form is always $\begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$; I is r by r .

Problem Set 3.4, page 163

- 1 $\begin{bmatrix} 2 & 4 & 6 & 4 & \mathbf{b}_1 \\ 2 & 5 & 7 & 6 & \mathbf{b}_2 \\ 2 & 3 & 5 & 2 & \mathbf{b}_3 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 4 & 6 & 4 & \mathbf{b}_1 \\ 0 & 1 & 1 & 2 & \mathbf{b}_2 - \mathbf{b}_1 \\ 0 & -1 & -1 & -2 & \mathbf{b}_3 - \mathbf{b}_1 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 4 & 6 & 4 & \mathbf{b}_1 \\ 0 & 1 & 1 & 2 & \mathbf{b}_2 - \mathbf{b}_1 \\ 0 & 0 & 0 & 0 & \mathbf{b}_3 + \mathbf{b}_2 - 2\mathbf{b}_1 \end{bmatrix}$
 $A\mathbf{x} = \mathbf{b}$ has a solution when $b_3 + b_2 - 2b_1 = 0$; the column space contains all combinations of $(2, 2, 2)$ and $(4, 5, 3)$. **This is the plane** $b_3 + b_2 - 2b_1 = 0$ (!). The nullspace contains all combinations of $s_1 = (-1, -1, 1, 0)$ and $s_2 = (2, -2, 0, 1)$; $x_{\text{complete}} = x_p + c_1 s_1 + c_2 s_2$;

$$[R \ d] = \begin{bmatrix} 1 & 0 & 1 & -2 & 4 \\ 0 & 1 & 1 & 2 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \text{ gives the particular solution } x_p = (4, -1, 0, 0).$$

$$2 \begin{bmatrix} 2 & 1 & 3 & \mathbf{b}_1 \\ 6 & 3 & 9 & \mathbf{b}_2 \\ 4 & 2 & 6 & \mathbf{b}_3 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 3 & \mathbf{b}_1 \\ 0 & 0 & 0 & \mathbf{b}_2 - 3\mathbf{b}_1 \\ 0 & 0 & 0 & \mathbf{b}_3 - 2\mathbf{b}_1 \end{bmatrix} \quad \text{Then } [R \ \mathbf{d}] = \begin{bmatrix} 1 & 1/2 & 3/2 & \mathbf{5} \\ 0 & 0 & 0 & \mathbf{0} \\ 0 & 0 & 0 & \mathbf{0} \end{bmatrix}$$

$A\mathbf{x} = \mathbf{b}$ has a solution when $b_2 - 3b_1 = 0$ and $b_3 - 2b_1 = 0$; $C(A) =$ line through $(2, 6, 4)$ which is the intersection of the planes $b_2 - 3b_1 = 0$ and $b_3 - 2b_1 = 0$; the nullspace contains all combinations of $\mathbf{s}_1 = (-1/2, 1, 0)$ and $\mathbf{s}_2 = (-3/2, 0, 1)$; particular solution $\mathbf{x}_p = \mathbf{d} = (5, 0, 0)$ and complete solution $\mathbf{x}_p + c_1\mathbf{s}_1 + c_2\mathbf{s}_2$.

$$3 \mathbf{x}_{\text{complete}} = \begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} -3 \\ 1 \\ 0 \end{bmatrix}. \quad \text{The matrix is singular but the equations are still solvable; } \mathbf{b} \text{ is in the column space. Our particular solution has free variable } y = 0.$$

$$4 \mathbf{x}_{\text{complete}} = \mathbf{x}_p + \mathbf{x}_n = \left(\frac{1}{2}, 0, \frac{1}{2}, 0\right) + x_2(-3, 1, 0, 0) + x_4(0, 0, -2, 1).$$

$$5 \begin{bmatrix} 1 & 2 & -2 & b_1 \\ 2 & 5 & -4 & b_2 \\ 4 & 9 & -8 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & -2 & b_1 \\ 0 & 1 & 0 & b_2 - 2b_1 \\ 0 & 0 & 0 & b_3 - 2b_1 - b_2 \end{bmatrix} \quad \text{solvable if } b_3 - 2b_1 - b_2 = 0.$$

Back-substitution gives the particular solution to $A\mathbf{x} = \mathbf{b}$ and the special solution to

$$A\mathbf{x} = \mathbf{0}: \mathbf{x} = \begin{bmatrix} 5b_1 - 2b_2 \\ b_2 - 2b_1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}.$$

$$6 \text{ (a) Solvable if } b_2 = 2b_1 \text{ and } 3b_1 - 3b_3 + b_4 = 0. \text{ Then } \mathbf{x} = \begin{bmatrix} 5b_1 - 2b_3 \\ b_3 - 2b_1 \end{bmatrix} = \mathbf{x}_p$$

$$\text{(b) Solvable if } b_2 = 2b_1 \text{ and } 3b_1 - 3b_3 + b_4 = 0. \mathbf{x} = \begin{bmatrix} 5b_1 - 2b_3 \\ b_3 - 2b_1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix}.$$

$$7 \begin{bmatrix} 1 & 3 & 1 & b_1 \\ 3 & 8 & 2 & b_2 \\ 2 & 4 & 0 & b_3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 3 & 1 & b_2 \\ 0 & -1 & -1 & b_2 - 3b_1 \\ 0 & -2 & -2 & b_3 - 2b_1 \end{bmatrix} \quad \text{One more step gives } [0 \ 0 \ 0 \ 0] = \text{row } 3 - 2(\text{row } 2) + 4(\text{row } 1) \text{ provided } b_3 - 2b_2 + 4b_1 = 0.$$

8 (a) Every \mathbf{b} is in $C(A)$: independent rows, only the zero combination gives $\mathbf{0}$.

(b) We need $b_3 = 2b_2$, because $(\text{row } 3) - 2(\text{row } 2) = \mathbf{0}$.

$$9 L[U \ \mathbf{c}] = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 & 5 & b_1 \\ 0 & 0 & 2 & 2 & b_2 - 2b_1 \\ 0 & 0 & 0 & 0 & b_3 + b_2 - 5b_1 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 & 5 & b_1 \\ 2 & 4 & 8 & 12 & b_2 \\ 3 & 6 & 7 & 13 & b_3 \end{bmatrix} \\ = [A \ \mathbf{b}]; \text{ particular } \mathbf{x}_p = (-9, 0, 3, 0) \text{ means } -9(1, 2, 3) + 3(3, 8, 7) = (0, 6, -6). \\ \text{This is } A\mathbf{x}_p = \mathbf{b}.$$

$$10 \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 4 \end{bmatrix} \text{ has } \mathbf{x}_p = (2, 4, 0) \text{ and } \mathbf{x}_{\text{null}} = (c, c, c).$$

11 A 1 by 3 system has at least **two** free variables. But \mathbf{x}_{null} in Problem 10 only has **one**.

12 (a) $\mathbf{x}_1 - \mathbf{x}_2$ and $\mathbf{0}$ solve $A\mathbf{x} = \mathbf{0}$ (b) $A(2\mathbf{x}_1 - 2\mathbf{x}_2) = \mathbf{0}$, $A(2\mathbf{x}_1 - \mathbf{x}_2) = \mathbf{b}$

13 (a) The particular solution \mathbf{x}_p is always multiplied by 1 (b) Any solution can be \mathbf{x}_p

$$\text{(c) } \begin{bmatrix} 3 & 3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 6 \\ 6 \end{bmatrix}. \text{ Then } \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ is shorter (length } \sqrt{2}) \text{ than } \begin{bmatrix} 2 \\ 0 \end{bmatrix} \text{ (length } 2)$$

(d) The only "homogeneous" solution in the nullspace is $\mathbf{x}_n = \mathbf{0}$ when A is invertible.

- 14 If column 5 has no pivot, x_5 is a *free* variable. The zero vector is *not* the only solution to $A\mathbf{x} = \mathbf{0}$. If this system $A\mathbf{x} = \mathbf{b}$ has a solution, it has *infinitely many* solutions.
- 15 If row 3 of U has no pivot, that is a *zero row*. $U\mathbf{x} = \mathbf{c}$ is only solvable provided $c_3 = 0$. $A\mathbf{x} = \mathbf{b}$ *might not be solvable*, because U may have other zero rows needing more $c_i = 0$.
- 16 The largest rank is 3. Then there is a pivot in every *row*. The solution *always exists*. The column space is \mathbf{R}^3 . An example is $A = [I \ F]$ for any 3 by 2 matrix F .
- 17 The largest rank of a 6 by 4 matrix is 4. Then there is a pivot in every *column*. The solution is *unique*. The nullspace contains only the zero *vector*. An example is $A = R = [I \ F]$ for any 4 by 2 matrix F .
- 18 Rank = 2; rank = 3 unless $q = 2$ (then rank = 2). Transpose has the same rank!
- 19 Both matrices A have rank 2. Always $A^T A$ and AA^T have **the same rank** as A .
- 20 $A = LU = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 3 & 4 & 1 & 0 \\ 0 & -3 & 0 & 1 \end{bmatrix}$; $A = LU \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 2 & -2 & 3 \\ 0 & 0 & 11 & -5 \end{bmatrix}$.
- 21 (a) $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ 0 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} + z \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$ (b) $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ 0 \end{bmatrix} + z \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$. The second equation in part (b) removed one special solution.
- 22 If $A\mathbf{x}_1 = \mathbf{b}$ and also $A\mathbf{x}_2 = \mathbf{b}$ then we can add $\mathbf{x}_1 - \mathbf{x}_2$ to any solution of $A\mathbf{x} = \mathbf{B}$: the solution \mathbf{x} is not unique. But there will be **no solution** to $A\mathbf{x} = \mathbf{B}$ if \mathbf{B} is not in the column space.
- 23 For A , $q = 3$ gives rank 1, every other q gives rank 2. For B , $q = 6$ gives rank 1, every other q gives rank 2. These matrices cannot have rank 3.
- 24 (a) $\begin{bmatrix} 1 \\ 1 \end{bmatrix} [x] = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$ has 0 or 1 solutions, depending on \mathbf{b} (b) $\begin{bmatrix} 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = [b]$ has infinitely many solutions for every b (c) There are 0 or ∞ solutions when A has rank $r < m$ and $r < n$: the simplest example is a zero matrix. (d) *one* solution for all \mathbf{b} when A is square and invertible (like $A = I$).
- 25 (a) $r < m$, always $r \leq n$ (b) $r = m$, $r < n$ (c) $r < m$, $r = n$ (d) $r = m = n$.
- 26 $\begin{bmatrix} 2 & 4 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow R = \begin{bmatrix} \mathbf{1} & 0 & -2 \\ 0 & \mathbf{1} & 2 \\ 0 & 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 2 & 4 & 4 \\ 0 & 3 & 6 \\ 0 & 0 & 5 \end{bmatrix} \rightarrow R = I$.
- 27 If U has n pivots, then R has n pivots **equal to 1**. Zeros above and below those pivots make $R = I$.
- 28 $\begin{bmatrix} 1 & 2 & 3 & \mathbf{0} \\ 0 & 0 & 4 & \mathbf{0} \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 0 & \mathbf{0} \\ 0 & 0 & 1 & \mathbf{0} \end{bmatrix}$; $\mathbf{x}_n = \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$; $\begin{bmatrix} 1 & 2 & 3 & \mathbf{5} \\ 0 & 0 & 4 & \mathbf{8} \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 0 & -\mathbf{1} \\ 0 & 0 & 1 & \mathbf{2} \end{bmatrix}$.
Free $x_2 = 0$ gives $\mathbf{x}_p = (-1, 0, 2)$ because the pivot columns contain I .
- 29 $[R \ \mathbf{d}] = \begin{bmatrix} 1 & 0 & 0 & \mathbf{0} \\ 0 & 0 & 1 & \mathbf{0} \\ 0 & 0 & 0 & \mathbf{0} \end{bmatrix}$ leads to $\mathbf{x}_n = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$; $[R \ \mathbf{d}] = \begin{bmatrix} 1 & 0 & 0 & -\mathbf{1} \\ 0 & 0 & 1 & \mathbf{2} \\ 0 & 0 & 0 & \mathbf{5} \end{bmatrix}$:
no solution because of the 3rd equation

$$30 \begin{bmatrix} 1 & 0 & 2 & 3 & 2 \\ 1 & 3 & 2 & 0 & 5 \\ 2 & 0 & 4 & 9 & 10 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2 & 3 & 2 \\ 0 & 3 & 0 & -3 & 3 \\ 0 & 0 & 0 & 3 & 6 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 2 & 0 & -4 \\ 0 & 1 & 0 & 0 & 3 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix}; \begin{bmatrix} -4 \\ 3 \\ 0 \\ 2 \end{bmatrix}; x_n = x_3 \begin{bmatrix} -2 \\ 0 \\ 1 \\ 0 \end{bmatrix}.$$

31 For $A = \begin{bmatrix} 1 & 1 \\ 0 & 2 \\ 0 & 3 \end{bmatrix}$, the only solution to $A\mathbf{x} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ is $\mathbf{x} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. B cannot exist since 2 equations in 3 unknowns cannot have a unique solution.

32 $A = \begin{bmatrix} 1 & 3 & 1 \\ 1 & 2 & 3 \\ 2 & 4 & 6 \\ 1 & 1 & 5 \end{bmatrix}$ factors into $LU = \begin{bmatrix} 1 & & & \\ 1 & 1 & & \\ 2 & 2 & 1 & \\ 1 & 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ and the rank is $r = 2$. The special solution to $A\mathbf{x} = \mathbf{0}$ and $U\mathbf{x} = \mathbf{0}$ is $s = (-7, 2, 1)$. Since $\mathbf{b} = (1, 3, 6, 5)$ is also the last column of A , a particular solution to $A\mathbf{x} = \mathbf{b}$ is $(0, 0, 1)$ and the complete solution is $\mathbf{x} = (0, 0, 1) + c\mathbf{s}$. (Or use the particular solution $\mathbf{x}_p = (7, -2, 0)$ with free variable $x_3 = 0$.)

For $\mathbf{b} = (1, 0, 0, 0)$ elimination leads to $U\mathbf{x} = (1, -1, 0, 1)$ and the fourth equation is $0 = 1$. No solution for this \mathbf{b} .

33 If the complete solution to $A\mathbf{x} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$ is $\mathbf{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ c \end{bmatrix}$ then $A = \begin{bmatrix} 1 & 0 \\ 3 & 0 \end{bmatrix}$.

34 (a) If $s = (2, 3, 1, 0)$ is the only special solution to $A\mathbf{x} = \mathbf{0}$, the complete solution is $\mathbf{x} = c\mathbf{s}$ (line of solution!). The rank of A must be $4 - 1 = 3$.

(b) The fourth variable x_4 is *not free* in s , and R must be $\begin{bmatrix} 1 & 0 & -2 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$.

(c) $A\mathbf{x} = \mathbf{b}$ can be solve for all \mathbf{b} , because A and R have *full row rank* $r = 3$.

35 For the $-1, 2, -1$ matrix $K(9 \text{ by } 9)$ and constant right side $\mathbf{b} = (10, \dots, 10)$, the solution $\mathbf{x} = K^{-1}\mathbf{b} = (45, 80, 105, 120, 125, 120, 105, 80, 45)$ rises and falls along the parabola $x_i = 50i - 5i^2$. (A formula for K^{-1} is later in the text.)

36 If $A\mathbf{x} = \mathbf{b}$ and $C\mathbf{x} = \mathbf{b}$ have the same solutions, A and C have the same shape and the same nullspace (take $\mathbf{b} = \mathbf{0}$). If $\mathbf{b} = \text{column 1 of } A$, $\mathbf{x} = (1, 0, \dots, 0)$ solves $A\mathbf{x} = \mathbf{b}$ so it solves $C\mathbf{x} = \mathbf{b}$. Then A and C share column 1. Other columns too: $A = C$!

Problem Set 3.5, page 178

$$1 \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = 0 \text{ gives } c_3 = c_2 = c_1 = 0. \text{ So those 3 column vectors are}$$

independent. But $\begin{bmatrix} 1 & 1 & 1 & 2 \\ 0 & 1 & 1 & 3 \\ 0 & 0 & 1 & 4 \end{bmatrix} [\mathbf{c}] = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ is solved by $\mathbf{c} = (1, 1, -4, 1)$. Then $\mathbf{v}_1 + \mathbf{v}_2 - 4\mathbf{v}_3 + \mathbf{v}_4 = \mathbf{0}$ (dependent).

2 $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ are independent (the -1 's are in different positions). All six vectors are on the plane $(1, 1, 1, 1) \cdot \mathbf{v} = 0$ so no four of these six vectors can be independent.

- 3 If $a = 0$ then column 1 = $\mathbf{0}$; if $d = 0$ then $b(\text{column 1}) - a(\text{column 2}) = \mathbf{0}$; if $f = 0$ then all columns end in zero (they are all in the xy plane, they must be dependent).
- 4 $U\mathbf{x} = \begin{bmatrix} a & b & c \\ 0 & d & e \\ 0 & 0 & f \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ gives $z = 0$ then $y = 0$ then $x = 0$. A square triangular matrix has independent columns (invertible matrix) when its diagonal has no zeros.
- 5 (a) $\begin{bmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \\ 2 & 3 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 \\ 0 & -5 & -7 \\ 0 & -1 & -5 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 3 \\ 0 & -5 & -7 \\ 0 & 0 & -18/5 \end{bmatrix}$: invertible \Rightarrow independent columns.
- (b) $\begin{bmatrix} 1 & 2 & -3 \\ -3 & 1 & 2 \\ 2 & -3 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & -3 \\ 0 & 7 & -7 \\ 0 & -7 & 7 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & -3 \\ 0 & 7 & -7 \\ 0 & 0 & 0 \end{bmatrix}$; $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$, columns add to $\mathbf{0}$.
- 6 Columns 1, 2, 4 are independent. Also 1, 3, 4 and 2, 3, 4 and others (but not 1, 2, 3). Same column numbers (not same columns!) for A .
- 7 The sum $\mathbf{v}_1 - \mathbf{v}_2 + \mathbf{v}_3 = \mathbf{0}$ because $(\mathbf{w}_2 - \mathbf{w}_3) - (\mathbf{w}_1 - \mathbf{w}_3) + (\mathbf{w}_1 - \mathbf{w}_2) = \mathbf{0}$. So the difference are *dependent* and the difference matrix is singular: $A = \begin{bmatrix} 0 & 1 & -1 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{bmatrix}$.
- 8 If $c_1(\mathbf{w}_2 + \mathbf{w}_3) + c_2(\mathbf{w}_1 + \mathbf{w}_3) + c_3(\mathbf{w}_1 + \mathbf{w}_2) = \mathbf{0}$ then $(c_2 + c_3)\mathbf{w}_1 + (c_1 + c_3)\mathbf{w}_2 + (c_1 + c_2)\mathbf{w}_3 = \mathbf{0}$. Since the \mathbf{w} 's are independent, $c_2 + c_3 = c_1 + c_3 = c_1 + c_2 = 0$. The only solution is $c_1 = c_2 = c_3 = 0$. Only this combination of $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ gives $\mathbf{0}$.
- 9 (a) The four vectors in \mathbf{R}^3 are the columns of a 3 by 4 matrix A . There is a nonzero solution to $A\mathbf{x} = \mathbf{0}$ because there is at least one free variable (b) Two vectors are dependent if $[\mathbf{v}_1 \ \mathbf{v}_2]$ has rank 0 or 1. (OK to say "they are on the same line" or "one is a multiple of the other" but *not* " \mathbf{v}_2 is a multiple of \mathbf{v}_1 "—since \mathbf{v}_1 might be $\mathbf{0}$.) (c) A nontrivial combination of \mathbf{v}_1 and $\mathbf{0}$ gives $\mathbf{0}$: $0\mathbf{v}_1 + 3(0, 0, 0) = \mathbf{0}$.
- 10 The plane is the nullspace of $A = [1 \ 2 \ -3 \ -1]$. Three free variables give three solutions $(x, y, z, t) = (2, -1, 0, 0)$ and $(3, 0, 1, 0)$ and $(1, 0, 0, 1)$. Combinations of those special solutions give more solutions (all solutions).
- 11 (a) Line in \mathbf{R}^3 (b) Plane in \mathbf{R}^3 (c) All of \mathbf{R}^3 (d) All of \mathbf{R}^3 .
- 12 \mathbf{b} is in the column space when $A\mathbf{x} = \mathbf{b}$ has a solution; \mathbf{c} is in the row space when $A^T\mathbf{y} = \mathbf{c}$ has a solution. *False*. The zero vector is always in the row space.
- 13 The column space and row space of A and U all have the same dimension = 2. *The row spaces of A and U are the same*, because the rows of U are combinations of the rows of A (and vice versa!).
- 14 $\mathbf{v} = \frac{1}{2}(\mathbf{v} + \mathbf{w}) + \frac{1}{2}(\mathbf{v} - \mathbf{w})$ and $\mathbf{w} = \frac{1}{2}(\mathbf{v} + \mathbf{w}) - \frac{1}{2}(\mathbf{v} - \mathbf{w})$. The two pairs *span* the same space. They are a basis when \mathbf{v} and \mathbf{w} are *independent*.
- 15 The n independent vectors span a space of dimension n . They are a *basis* for that space. If they are the columns of A then m is *not less* than n ($m \geq n$).

16 These bases are not unique! (a) $(1, 1, 1, 1)$ for the space of all constant vectors (c, c, c, c) (b) $(1, -1, 0, 0), (1, 0, -1, 0), (1, 0, 0, -1)$ for the space of vectors with sum of components = 0 (c) $(1, -1, -1, 0), (1, -1, 0, -1)$ for the space perpendicular to $(1, 1, 0, 0)$ and $(1, 0, 1, 1)$ (d) The columns of I are a basis for its column space, the empty set is a basis (by convention) for $N(I) = \{\text{zero vector}\}$.

17 The column space of $U = \begin{bmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \end{bmatrix}$ is \mathbf{R}^2 so take any bases for \mathbf{R}^2 ; (row 1 and row 2) or (row 1 and row 1 + row 2) and (row 1 and - row 2) are bases for the row spaces of U .

18 (a) The 6 vectors *might not* span \mathbf{R}^4 (b) The 6 vectors *are not* independent (c) Any four *might be* a basis.

19 n -independent columns \Rightarrow rank n . Columns span $\mathbf{R}^m \Rightarrow$ rank m . Columns are basis for $\mathbf{R}^m \Rightarrow$ rank = $m = n$. The rank counts the number of *independent* columns.

20 One basis is $(2, 1, 0), (-3, 0, 1)$. A basis for the intersection with the xy plane is $(2, 1, 0)$. The normal vector $(1, -2, 3)$ is a basis for the line perpendicular to the plane.

21 (a) The only solution to $A\mathbf{x} = \mathbf{0}$ is $\mathbf{x} = \mathbf{0}$ because *the columns are independent* (b) $A\mathbf{x} = \mathbf{b}$ is solvable because *the columns span \mathbf{R}^5* . Key point: A basis gives exactly one solution for every \mathbf{b} .

22 (a) True (b) False because the basis vectors for \mathbf{R}^6 might not be in \mathbf{S} .

23 Columns 1 and 2 are bases for the (**different**) column spaces of A and U ; rows 1 and 2 are bases for the (**equal**) row spaces of A and U ; $(1, -1, 1)$ is a basis for the (**equal**) nullspaces.

24 (a) *False* $A = \begin{bmatrix} 1 & 1 \end{bmatrix}$ has dependent columns, independent row (b) *False* column space \neq row space for $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ (c) *True*: Both dimensions = 2 if A is invertible, dimensions = 0 if $A = 0$, otherwise dimensions = 1 (d) *False*, columns may be dependent, in that case not a basis for $\mathcal{C}(A)$.

25 A has rank 2 if $c = 0$ and $d = 2$; $B = \begin{bmatrix} c & d \\ d & c \end{bmatrix}$ has rank 2 except when $c = d$ or $c = -d$.

26 (a) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

(b) Add $\begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$

(c) $\begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}$.

These are simple bases (among many others) for (a) diagonal matrices (b) symmetric matrices (c) skew-symmetric matrices. The dimensions are 3, 6, 3.

- 27 $I, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$; echelon matrices do *not* form a subspace; they *span* the upper triangular matrices (not every U is echelon).
- 28 $\begin{bmatrix} 1 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & -1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & -1 \end{bmatrix}; \begin{bmatrix} 1 & -1 & 0 \\ -1 & 1 & 0 \end{bmatrix}$ and $\begin{bmatrix} 1 & 0 & -1 \\ -1 & 0 & 1 \end{bmatrix}$.
- 29 (a) The invertible matrices span the space of all 3 by 3 matrices (b) The rank one matrices also span the space of all 3 by 3 matrices (c) I by itself spans the space of all multiples cI .
- 30 $\begin{bmatrix} -1 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} -1 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ -1 & 2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ -1 & 0 & 2 \end{bmatrix}$.
- 31 (a) $y(x) = \text{constant } C$ (b) $y(x) = 3x$ this is one basis for the 2 by 3 matrices with $(2, 1, 1)$ in their nullspace (4-dim subspace). (c) $y(x) = 3x + C = y_p + y_n$ solves $dy/dx = 3$.
- 32 $y(0) = 0$ requires $A + B + C = 0$. One basis is $\cos x - \cos 2x$ and $\cos x - \cos 3x$.
- 33 (a) $y(x) = e^{2x}$ is a basis for, all solutions to $y' = 2y$ (b) $y = x$ is a basis for all solutions to $dy/dx = y/x$ (First-order linear equation \Rightarrow 1 basis function in solution space).
- 34 $y_1(x), y_2(x), y_3(x)$ can be $x, 2x, 3x$ (dim 1) or $x, 2x, x^2$ (dim 2) or x, x^2, x^3 (dim 3).
- 35 Basis $1, x, x^2, x^3$, for cubic polynomials; basis $x - 1, x^2 - 1, x^3 - 1$ for the subspace with $p(1) = 0$.
- 36 Basis for \mathbf{S} : $(1, 0, -1, 0), (0, 1, 0, 0), (1, 0, 0, -1)$; basis for \mathbf{T} : $(1, -1, 0, 0)$ and $(0, 0, 2, 1)$; $\mathbf{S} \cap \mathbf{T} =$ multiples of $(3, -3, 2, 1) =$ nullspace for 3 equation in \mathbf{R}^4 has dimension 1.
- 37 The subspace of matrices that have $AS = SA$ has dimension *three*.
- 38 (a) No, 2 vectors don't span \mathbf{R}^3 (b) No, 4 vectors in \mathbf{R}^3 are dependent (c) Yes, a basis (d) No, these three vectors are dependent
- 39 If the 5 by 5 matrix $[A \ \mathbf{b}]$ is invertible, \mathbf{b} is not a combination of the columns of A . If $[A \ \mathbf{b}]$ is singular, and the 4 columns of A are independent, \mathbf{b} is a combination of those columns. In this case $A\mathbf{x} = \mathbf{b}$ has a solution.
- 40 (a) The functions $y = \sin x, y = \cos x, y = e^x, y = e^{-x}$ are a basis for solutions to $d^4y/dx^4 = y(x)$.
 (b) A particular solution to $d^4y/dx^4 = y(x) + 1$ is $y(x) = -1$. The complete solution is $y(x) = -1 + c_1 \sin x + c_2 \cos x + c_3 e^x + c_4 e^{-x}$ (or use another basis for the nullspace of the 4th derivative).
- 41 $I = \begin{bmatrix} & 1 & & & & \\ 1 & & & & & \\ & & 1 & & & \\ & & & 1 & & \\ & & & & 1 & \\ & & & & & 1 \end{bmatrix} - \begin{bmatrix} & 1 & & & & \\ & & 1 & & & \\ 1 & & & & & \\ & & & 1 & & \\ & & & & 1 & \\ & & & & & 1 \end{bmatrix} + \begin{bmatrix} & & & 1 & & \\ & & & & 1 & \\ & & & 1 & & \\ & & & & 1 & \\ & & & & & 1 \end{bmatrix} + \begin{bmatrix} & & & & 1 & \\ & & & & & 1 \\ & & & & 1 & \\ & & & & & 1 \\ & & & & & & 1 \end{bmatrix} - \begin{bmatrix} & & & & & 1 \\ & & & & & & 1 \\ & & & & & 1 & \\ & & & & & & 1 \\ & & & & & & & 1 \end{bmatrix}$. The six P 's are dependent.
 Those five are independent: The 4th has $P_{11} = 1$ and cannot be a combination of the others. Then the 2nd cannot be (from $P_{32} = 1$) and also 5th ($P_{32} = 1$). Continuing, a nonzero combination of all five could not be zero. Further challenge: How many independent 4 by 4 permutation matrices?

- 42 The dimension of \mathcal{S} spanned by all rearrangements of \mathbf{x} is (a) zero when $\mathbf{x} = \mathbf{0}$ (b) one when $\mathbf{x} = (1, 1, 1, 1)$ (c) three when $\mathbf{x} = (1, 1, -1, -1)$ because all rearrangements of this \mathbf{x} are perpendicular to $(1, 1, 1, 1)$ (d) four when the \mathbf{x} 's are not equal and don't add to zero. **No \mathbf{x} gives $\dim \mathcal{S} = 2$.** I owe this nice problem to Mike Artin—the answers are the same in higher dimensions: $0, 1, n - 1, n$.
- 43 The problem is to show that the \mathbf{u} 's, \mathbf{v} 's, \mathbf{w} 's together are independent. We know the \mathbf{u} 's and \mathbf{v} 's together are a basis for V , and the \mathbf{u} 's and \mathbf{w} 's together are a basis for W . Suppose a combination of \mathbf{u} 's, \mathbf{v} 's, \mathbf{w} 's gives $\mathbf{0}$. **To be proved:** All coefficients = zero.
Key idea: In that combination giving $\mathbf{0}$, the part \mathbf{x} from the \mathbf{u} 's and \mathbf{v} 's is in V . So the part from the \mathbf{w} 's is $-\mathbf{x}$. This part is now in V and also in W . But if $-\mathbf{x}$ is in $V \cap W$ it is a combination of \mathbf{u} 's only. Now the combination uses only \mathbf{u} 's and \mathbf{v} 's (independent in V !) so all coefficients of \mathbf{u} 's and \mathbf{v} 's must be zero. Then $\mathbf{x} = \mathbf{0}$ and the coefficients of the \mathbf{w} 's are also zero.
- 44 The inputs to an m by n matrix fill \mathbf{R}^n . The outputs (column space!) have dimension r . The nullspace has $n - r$ special solutions. The formula becomes $r + (n - r) = n$.
- 45 If the left side of $\dim(V) + \dim(W) = \dim(V \cap W) + \dim(V + W)$ is greater than n , then $\dim(V \cap W)$ must be greater than zero. So $V \cap W$ contains nonzero vectors.
- 46 If $A^2 =$ zero matrix, this says that each column of A is in the nullspace of A . If the column space has dimension r , the nullspace has dimension $10 - r$, and we must have $r \leq 10 - r$ and $r \leq 5$.

Problem Set 3.6, page 190

- 1 (a) Row and column space dimensions = 5, nullspace dimension = 4, $\dim(N(A^T)) = 2$ sum = $16 = m + n$ (b) Column space is \mathbf{R}^3 ; left nullspace contains only $\mathbf{0}$.
- 2 A : Row space basis = row 1 = $(1, 2, 4)$; nullspace $(-2, 1, 0)$ and $(-4, 0, 1)$; column space basis = column 1 = $(1, 2)$; left nullspace $(-2, 1)$. B : Row space basis = both rows = $(1, 2, 4)$ and $(2, 5, 8)$; column space basis = two columns = $(1, 2)$ and $(2, 5)$; nullspace $(-4, 0, 1)$; left nullspace basis is empty because the space contains only $\mathbf{y} = \mathbf{0}$.
- 3 Row space basis = rows of $U = (0, 1, 2, 3, 4)$ and $(0, 0, 0, 1, 2)$; column space basis = pivot columns (of A not U) = $(1, 1, 0)$ and $(3, 4, 1)$; nullspace basis $(1, 0, 0, 0, 0)$, $(0, 2, -1, 0, 0)$, $(0, 2, 0, -2, 1)$; left nullspace $(1, -1, 1) =$ last row of E^{-1} !
- 4 (a) $\begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$ (b) Impossible: $r + (n - r)$ must be 3 (c) $[1 \ 1]$ (d) $\begin{bmatrix} -9 & -3 \\ 3 & 1 \end{bmatrix}$
 (e) *Impossible* Row space = column space requires $m = n$. Then $m - r = n - r$; nullspaces have the same dimension. Section 4.1 will prove $N(A)$ and $N(A^T)$ orthogonal to the row and column spaces respectively—here those are the same space.
- 5 $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 1 & 0 \end{bmatrix}$ has those rows spanning its row space $B = [1 \ -2 \ 1]$ has the same rows spanning its nullspace and $BA^T = 0$.
- 6 A : dim **2, 2, 2, 1**: Rows $(0, 3, 3, 3)$ and $(0, 1, 0, 1)$; columns $(3, 0, 1)$ and $(3, 0, 0)$; nullspace $(1, 0, 0, 0)$ and $(0, -1, 0, 1)$; $N(A^T)$ $(0, 1, 0)$. B : dim **1, 1, 0, 2** Row space (1) , column space $(1, 4, 5)$, nullspace: empty basis, $N(A^T)$ $(-4, 1, 0)$ and $(-5, 0, 1)$.

- 7 Invertible 3 by 3 matrix A : row space basis = column space basis = $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$; nullspace basis and left nullspace basis are *empty*. Matrix $B = \begin{bmatrix} A & A \end{bmatrix}$: row space basis $(1, 0, 0, 1, 0, 0)$, $(0, 1, 0, 0, 1, 0)$ and $(0, 0, 1, 0, 0, 1)$; column space basis $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$; nullspace basis $(-1, 0, 0, 1, 0, 0)$ and $(0, -1, 0, 0, 1, 0)$ and $(0, 0, -1, 0, 0, 1)$; left nullspace basis is empty.
- 8 $\begin{bmatrix} I & 0 \end{bmatrix}$ and $\begin{bmatrix} I & I; & 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \end{bmatrix}$ = 3 by 2 have row space dimensions = 3, 3, 0 = column space dimensions; nullspace dimensions 2, 3, 2; left nullspace dimensions 0, 2, 3.
- 9 (a) Same row space and nullspace. So rank (dimension of row space) is the same
 (b) Same column space and left nullspace. Same rank (dimension of column space).
- 10 For **rand** (3), almost surely rank = 3, nullspace and left nullspace contain only $(0, 0, 0)$.
 For **rand** (3, 5) the rank is almost surely 3 and the dimension of the nullspace is 2.
- 11 (a) No solution means that $r < m$. Always $r \leq n$. Can't compare m and n here.
 (b) Since $m - r > 0$, the left nullspace must contain a nonzero vector.
- 12 A neat choice is $\begin{bmatrix} 1 & 1 \\ 0 & 2 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 1 \\ 1 & 2 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 2 & 1 \\ 2 & 4 & 0 \\ 1 & 0 & 1 \end{bmatrix}$; $r + (n - r) = n = 3$ does not match $2 + 2 = 4$. Only $\mathbf{v} = \mathbf{0}$ is in both $N(A)$ and $C(A^T)$.
- 13 (a) *False*: Usually row space \neq column space (same dimension!) (b) *True*: A and $-A$ have the same four subspaces (c) *False* (choose A and B same size and invertible: then they have the same four subspaces)
- 14 Row space basis can be the nonzero rows of U : $(1, 2, 3, 4)$, $(0, 1, 2, 3)$, $(0, 0, 1, 2)$; nullspace basis $(0, 1, -2, 1)$ as for U ; column space basis $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$ (happen to have $C(A) = C(U) = \mathbf{R}^3$); left nullspace has empty basis.
- 15 After a row exchange, the row space and nullspace stay the same; $(2, 1, 3, 4)$ is in the new left nullspace after the row exchange.
- 16 If $A\mathbf{v} = \mathbf{0}$ and \mathbf{v} is a row of A then $\mathbf{v} \cdot \mathbf{v} = 0$.
- 17 Row space = yz plane; column space = xy plane; nullspace = x axis; left nullspace = z axis. For $I + A$: Row space = column space = \mathbf{R}^3 , both nullspaces contain only the zero vector.
- 18 Row $3 - 2$ row $2 +$ row $1 =$ zero row so the vectors $c(1, -2, 1)$ are in the left nullspace. The same vectors happen to be in the nullspace (an accident for this matrix).
- 19 (a) Elimination on $A\mathbf{x} = \mathbf{0}$ leads to $0 = b_3 - b_2 - b_1$ so $(-1, -1, 1)$ is in the left nullspace. (b) 4 by 3: Elimination leads to $b_3 - 2b_1 = 0$ and $b_4 + b_2 - 4b_1 = 0$, so $(-2, 0, 1, 0)$ and $(-4, 1, 0, 1)$ are in the left nullspace. *Why?* Those vectors multiply the matrix to give *zero rows*. Section 4.1 will show another approach: $A\mathbf{x} = \mathbf{b}$ is solvable (\mathbf{b} is in $C(A)$) when \mathbf{b} is orthogonal to the left nullspace.
- 20 (a) Special solutions $(-1, 2, 0, 0)$ and $(-\frac{1}{4}, 0, -3, 1)$ are perpendicular to the rows of R (and then ER). (b) $A^T\mathbf{y} = \mathbf{0}$ has 1 independent solution = last row of E^{-1} . ($E^{-1}A = R$ has a zero row, which is just the transpose of $A^T\mathbf{y} = \mathbf{0}$).
- 21 (a) \mathbf{u} and \mathbf{w} (b) \mathbf{v} and \mathbf{z} (c) rank < 2 if \mathbf{u} and \mathbf{w} are dependent or if \mathbf{v} and \mathbf{z} are dependent (d) The rank of $\mathbf{u}\mathbf{v}^T + \mathbf{w}\mathbf{z}^T$ is 2.
- 22 $A = \begin{bmatrix} \mathbf{u} & \mathbf{w} \end{bmatrix} \begin{bmatrix} \mathbf{v}^T & \mathbf{z}^T \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 2 & 2 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 4 & 2 \\ 5 & 1 \end{bmatrix}$ has column space spanned by \mathbf{u} and \mathbf{w} , row space spanned by \mathbf{v} and \mathbf{z} .

- 23 As in Problem 22: Row space basis $(3, 0, 3), (1, 1, 2)$; column space basis $(1, 4, 2), (2, 5, 7)$; the rank of $(3 \text{ by } 2)$ times $(2 \text{ by } 3)$ cannot be larger than the rank of either factor, so $\text{rank} \leq 2$ and the $3 \text{ by } 3$ product is not invertible.
- 24 $A^T \mathbf{y} = \mathbf{d}$ puts \mathbf{d} in the row space of A ; unique solution if the left nullspace (nullspace of A^T) contains only $\mathbf{y} = \mathbf{0}$.
- 25 (a) True (A and A^T have the same rank) (b) False $A = [1 \ 0]$ and A^T have very different left nullspaces (c) False (A can be invertible and unsymmetric even if $C(A) = C(A^T)$) (d) True (The subspaces for A and $-A$ are always the same. If $A^T = A$ or $A^T = -A$ they are also the same for A^T)
- 26 The rows of $C = AB$ are combinations of the rows of B . So $\text{rank } C \leq \text{rank } B$. Also $\text{rank } C \leq \text{rank } A$, because the columns of C are combinations of the columns of A .
- 27 Choose $d = bc/a$ to make $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ a rank-1 matrix. Then the row space has basis (a, b) and the nullspace has basis $(-b, a)$. Those two vectors are perpendicular!
- 28 B and C (checkers and chess) both have rank 2 if $p \neq 0$. Row 1 and 2 are a basis for the row space of C , $B^T \mathbf{y} = \mathbf{0}$ has 6 special solutions with -1 and 1 separated by a zero; $N(C^T)$ has $(-1, 0, 0, 0, 0, 0, 1)$ and $(0, -1, 0, 0, 0, 0, 1, 0)$ and columns 3, 4, 5, 6 of I ; $N(C)$ is a challenge.
- 29 $a_{11} = 1, a_{12} = 0, a_{13} = 1, a_{22} = 0, a_{32} = 1, a_{31} = 0, a_{23} = 1, a_{33} = 0, a_{21} = 1$.
- 30 The subspaces for $A = \mathbf{u}\mathbf{v}^T$ are pairs of orthogonal lines (\mathbf{v} and \mathbf{v}^\perp , \mathbf{u} and \mathbf{u}^\perp). If B has those same four subspaces then $B = cA$ with $c \neq 0$.
- 31 (a) $AX = 0$ if each column of X is a multiple of $(1, 1, 1)$; $\dim(\text{nullspace}) = 3$. (b) If $AX = B$ then all columns of B add to zero; dimension of the B 's = 6. (c) $3 + 6 = \dim(M^{3 \times 3}) = 9$ entries in a $3 \text{ by } 3$ matrix.
- 32 The key is equal row spaces. First row of $A =$ combination of the rows of B : only possible combination (notice I) is 1 (row 1 of B). Same for each row so $F = G$.

Problem Set 4.1, page 202

- 1 Both nullspace vectors are orthogonal to the row space vector in \mathbf{R}^3 . The column space is perpendicular to the nullspace of A^T (two lines in \mathbf{R}^2 because $\text{rank} = 1$).
- 2 The nullspace of a $3 \text{ by } 2$ matrix with rank 2 is \mathbf{Z} (only zero vector) so $\mathbf{x}_n = \mathbf{0}$, and row space = \mathbf{R}^2 . Column space = plane perpendicular to left nullspace = line in \mathbf{R}^3 .
- 3 (a) $\begin{bmatrix} 1 & 2 & -3 \\ 2 & -3 & 1 \\ -3 & 5 & -2 \end{bmatrix}$ (b) Impossible, $\begin{bmatrix} 2 \\ -3 \\ 5 \end{bmatrix}$ not orthogonal to $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ (c) $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ in $C(A)$ and $N(A^T)$ is impossible: not perpendicular (d) Need $A^2 = 0$; take $A = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$ (e) $(1, 1, 1)$ in the nullspace (columns add to $\mathbf{0}$) and also row space; no such matrix.
- 4 If $AB = 0$, the columns of B are in the nullspace of A . The rows of A are in the left nullspace of B . If $\text{rank} = 2$, those four subspaces would have dimension 2 which is impossible for $3 \text{ by } 3$.
- 5 (a) If $A\mathbf{x} = \mathbf{b}$ has a solution and $A^T \mathbf{y} = \mathbf{0}$, then \mathbf{y} is perpendicular to \mathbf{b} . $\mathbf{b}^T \mathbf{y} = (A\mathbf{x})^T \mathbf{y} = \mathbf{x}^T (A^T \mathbf{y}) = 0$. (b) If $A^T \mathbf{y} = (1, 1, 1)$ has a solution, $(1, 1, 1)$ is in the row space and is orthogonal to every \mathbf{x} in the nullspace.