

$$\begin{aligned}
 yb + (\lambda - \mu)\beta &= \bar{y}b + \bar{\lambda}\beta - \lceil \mu^* \rceil \beta \\
 &\leq y^*b + (\lambda^* + \lceil \mu^* \rceil - \mu^*)\beta - \lceil \mu^* \rceil \beta
 \end{aligned}$$

since $(y^*, \lambda^* + \lceil \mu^* \rceil - \mu^*)$ is feasible for the minimum in (5.3), and $(\bar{y}, \bar{\lambda})$ is an optimum solution. We conclude that

$$yb + (\lambda - \mu)\beta \leq y^*b + (\lambda^* - \mu^*)\beta,$$

proving that (y, λ, μ) is an integral optimum solution for the minimum in (5.2). □

The following statements are straightforward consequences of the definition of TDI-systems: A system $Ax = b, x \geq 0$ is TDI if $\min\{yb : yA \geq c\}$ has an integral optimum solution y for each integral vector c for which the minimum is finite. A system $Ax \leq b, x \geq 0$ is TDI if $\min\{yb : yA \geq c, y \geq 0\}$ has an integral optimum solution y for each integral vector c for which the minimum is finite. One may ask whether there are matrices A such that $Ax \leq b, x \geq 0$ is TDI for each integral vector b . It will turn out that these matrices are exactly the totally unimodular matrices.

5.4 Totally Unimodular Matrices

Definition 5.18. A matrix A is totally unimodular if each subdeterminant of A is 0, +1, or -1.

In particular, each entry of a totally unimodular matrix must be 0, +1, or -1. The main result of this section is:

Theorem 5.19. (Hoffman and Kruskal [1956]) An integral matrix A is totally unimodular if and only if the polyhedron $\{x : Ax \leq b, x \geq 0\}$ is integral for each integral vector b .

Proof: Let A be an $m \times n$ -matrix and $P := \{x : Ax \leq b, x \geq 0\}$. Observe that the minimal faces of P are vertices.

To prove necessity, suppose that A is totally unimodular. Let b be some integral vector and x a vertex of P . x is the solution of $A'x = b'$ for some subsystem

$$A'x \leq b' \text{ of } \begin{pmatrix} A \\ -I \end{pmatrix} x \leq \begin{pmatrix} b \\ 0 \end{pmatrix}, \text{ with } A' \text{ being a nonsingular } n \times n\text{-matrix.}$$

Since A is totally unimodular, $|\det A'| = 1$, so by Cramer's rule $x = (A')^{-1}b'$ is integral.

We now prove sufficiency. Suppose that the vertices of P are integral for each integral vector b . Let A' be some nonsingular $k \times k$ -submatrix of A . We have to show $|\det A'| = 1$. W.l.o.g., A' contains the elements of the first k rows and columns of A .




Fig. 5.2.

Consider the integral $m \times m$ -matrix B consisting of the first k and the last $m - k$ columns of $(A \ I)$ (see Figure 5.2). Obviously, $|\det B| = |\det A'|$.

To prove $|\det B| = 1$, we shall prove that B^{-1} is integral. Since $\det B \det B^{-1} = 1$, this implies that $|\det B| = 1$, and we are done.

Let $i \in \{1, \dots, m\}$; we prove that $B^{-1}e_i$ is integral. Choose an integral vector y such that $z := y + B^{-1}e_i \geq 0$. Then $b := Bz = By + e_i$ is integral. We add zero components to z in order to obtain z' with

$$(A \ I)z' = Bz = b.$$

Now z'' , consisting of the first n components of z' , belongs to P . Furthermore, n linearly independent constraints are satisfied with equality, namely the first k and the last $n - k$ inequalities of $\begin{pmatrix} A \\ -I \end{pmatrix} z'' \leq 0$. Hence z'' is a vertex of P .

By our assumption z'' is integral. But then z' must also be integral: its first n components are the components of z'' , and the last m components are the slack variables $b - Az''$ (and A and b are integral). So z is also integral, and hence $B^{-1}e_i = z - y$ is integral. \square

The above proof is due to Veinott and Dantzig [1968].

Corollary 5.20. *An integral matrix A is totally unimodular if and only if for all integral vectors b and c both optima in the LP duality equation*

$$\max \{cx : Ax \leq b, x \geq 0\} = \min \{yb : y \geq 0, yA \geq c\}$$

are attained by integral vectors (if they are finite).

Proof: This follows from that the transpose of a tota

Let us reformulate thes

Corollary 5.21. *An integ system $Ax \leq b, x \geq 0$ is T*

Proof: If A (and thus A^T) Theorem $\min \{yb : yA \geq$ vector b and each integral words, the system $Ax \leq b,$

To show the converse, s b . Then by Corollary 5.14, each integral vector b . By

This is not the only wa a certain system is TDI. Th this will be used several tin

Lemma 5.22. *Let $Ax \leq b, b \in \mathbb{R}^m$. Suppose that for ec optimum solution, it has onu components of y^* form a to*

Proof: Let $c \in \mathbb{Z}^n$, and let 0) such that the rows of A totally unimodular matrix A

$$\min \{yb : yA \geq$$

where b' consists of the con the inequality " \leq " of (5.4), the LP on the left-hand side follows from the fact that y the LP on the right-hand sid

Since A' is totally unim optimum solution (by the Ho with zeros we obtain an intu completing the proof.

A very useful criterion f

Theorem 5.23. (Ghouila-H unimodular if and only if for R_2 such that