

Lab 03Name: [WebCT Administrator \(Preview\)](#)Start time: [October 27, 2003 11:45am](#) : Number of questions: 8[Finish](#)[Help](#)

This set of questions goes with the pages of applets and activities for [Lab 03](#). Use the applets and activities there to answer the questions.

Question 1 (1 point)

Refer to lab page 2. The initial measure of angle CBA is 55.8° . *Use the lengths of sides in the figure* and the definition of the cosecant function as a ratio to calculate $\csc(55.8^\circ)$. Report your answer correct to 3 decimal places.

Answer

[Save answer](#)**Question 2 (1 point)**

Refer to lab page 2. How long is the hypotenuse of an isosceles right triangle whose sides are of length 3 units? Drag the points A and B to decide. Answer correct to 2 decimal places.

Answer

[Save answer](#)**Question 3 (1 point)**

Refer to lab page 2. Suppose we want the measure in degrees of the smallest angle in a 5-12-13 right triangle. The units in the figure aren't the right size to make a picture easy to interpret. Figure out a way to use similar triangles in the figure so you can view this angle in a triangle that fits in the figure's viewing area. Give the number of degrees correct to the nearest tenth of a degree.

Answer

Save answer

Question 4 (1 point)

Refer to lab page 2. "Solving" a right triangle involves giving all angle measures and side lengths, once you are given some partial information. Eventually we will learn how to solve an arbitrary triangle analytically, but with the figure above you can recover missing information for the right triangle directly, by adjusting the positions of the points A , B , and C . Adjust the figure to find the length of the hypotenuse if AC (side b in the figure) is 2.75 inches and angle CBA is 50.0° . Report the length to as many decimal places as the figure provides.

Answer

Save answer

Question 5 (1 point)

Refer to lab page 2. "Solving" a right triangle involves giving all angle measures and side lengths, once you are given some partial information. Eventually we will learn how to solve an arbitrary triangle analytically, but with the figure above you can recover missing information for the right triangle directly, by adjusting the positions of the points A , B , and C . Adjust the figure to make AC (side b in the figure) 2.75 inches and angle CBA 50.0° . Find the measure of the other acute angle in degrees.

Answer

Save answer

Question 6 (3 points)

Refer to lab page 2. How about the tangent of 90° ? What problem arises with the ratio defining the tangent function when angle CBA gets larger and larger, approaching 90° ? Explain the situation in a sentence.

Refer to lab page 3. Graph the function $\tan(x)$. How does the problem above show up as a feature of the function graph? Explain the situation in a sentence.

Equation Create new equation ▼ Equation editor

Save answer

Question 7 (1 point)

Refer to lab page 2. Study limiting ratios in the figure to decide what value should be assigned to $\sec(0^\circ)$.

Answer

Save answer

Question 8 (1 point)

Use the Windows calculator to answer this question. Besides degrees and radians, another measure of angle size that is sometimes used is the *grad*. Scientific calculators often allow you to specify *deg*, *rad*, or *grad* as a mode for doing trigonometric calculations. Perform the following experiment to figure out the relationship between the different ways of measuring angles. Pick a convenient angle in degrees (say 45) and calculate the sin of the angle. Now set the calculator to grad mode and find the smallest positive grad measure of the angle that works to give a matching value of the sin function. Scale your answer up to decide how many grads there are in a complete revolution, corresponding to 360° .

Answer

Save answer

Finish

Help